

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for AM31179AF-N

### IL13 receptor alpha 2 (IL13RA2) Mouse Monoclonal Antibody [Clone ID: B-D13]

### **Product data:**


Product Type:	Primary Antibodies
Clone Name:	B-D13
Applications:	FC, IHC
Recommended Dilution:	Flow Cytometry. Immunohistochemistry on Paraffin Embedded Sections. Immunohistochemistry on Frozen Sections and methanol fixed cells.
Reactivity:	Human
Host:	Mouse
lsotype:	IgG1
Clonality:	Monoclonal
Immunogen:	IL-13Ra2 transfected CHO cell line. <b>Hybridoma:</b> Myeloma X63/AG.8653 x Balb/c spleen cells.
Specificity:	Recognizes the soluble antigen. Recognizes the IL-13 receptor a2 (IL-13 Ra2), a 44 kDa protein.
Formulation:	Phosphate-buffered saline without carrier and preservatives. This prodcut is sterile-filtered through 0.22 µm and treated to remove endotoxins. State: Azide Free State: Liquid purified IgG fraction
Concentration:	lot specific
Purification:	Ion Exchange Chromatography
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C.
Stability:	Shelf life: one year from despatch.
Gene Name:	interleukin 13 receptor subunit alpha 2
Database Link:	<u>Entrez Gene 3598 Human</u> <u>Q14627</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	IL13 receptor alpha 2 (IL13RA2) Mouse Monoclonal Antibody [Clone ID: B-D13] – AM31179AF-N
Background:	IL-13 receptor alpha 2 belongs to the type I cytokine receptor family; Contains 1 fibronectin type-III domain; Binds as a monomer with high affinity to interleukin-13 (IL13), but not to interleukin-4 (IL4). The WSXWS motif appears to be necessary for proper protein folding and thereby efficient intracellular transport and cell-surface receptor binding. The box 1 motif is required for JAK interaction and/or activation.
Synonyms:	Interleukin-13 receptor alpha-2 chain, IL13RA2, IL-13RA-2, IL13R, IL13 receptor alpha-2, IL- 13R-alpha-2
Protein Families	: Druggable Genome, Transmembrane
Protein Pathway	s: Jak-STAT signaling pathway

## Product images:



A typical staining pattern with the B-D13 monoclonal antibody on A375 cell line.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US