

Product datasheet for AM50239PU-S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD56 (NCAM1) Mouse Monoclonal Antibody [Clone ID: 123A8; same 56C04]

Product data:

Product Type: Primary Antibodies

Clone Name: 123A8; same 56C04

Applications: FC, IF, IHC, IP, WB

Recommended Dilution: ELISA: Use BSA free Antibody for Coating.

Western Blot (Not tested): 0.5-1 μg/ml.

Immunoprecipitation (Not tested): 1-2 μg/500 μg protein lysate.

Flow Cytometry (Not tested): 0.5-1 μ g/106 cells.

Immunofluorescence: 1-2 µg/ml.

Immunohistochemistry on Frozen and Formalin-Fixed Paraffin Sections: 0.5-1 µg/ml for

30 minutes at RT.

Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH

6.0, for 10-20 min followed by cooling at RT for 20 minutes. *Positive Control*: Cerebellum, Pancreas, Neuroblastoma.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Membrane preparation of a small cell lung carcinoma.

Specificity: This Monoclonal Antibody reacts with an extracellular domain (close to transmembrane) of

CD56/NCAM. Three isoforms of neural cell adhesion molecule (NCAM) are produced by differential splicing of the RNA transcript from a single gene. The 135kDa isoform is the basic

molecule, which is glycosylated or sialylated to produce the mature species.

Anti-CD56 recognizes two proteins of the neural cell adhesion molecule, the basic molecule expressed on most neuroectodermally derived tissues and neoplasms (e.g. retinoblastoma, medulloblastomas, astrocytomas, neuroblastomas, and small cell carcinomas). It is also

expressed on some mesodermally derived tumors (rhabdomyosarcoma).

Anti-CD56 plays an important role in the diagnosis of nodal and nasal NK/T-cell lymphomas.

Cellular Localization: Cell surface.





Formulation: 10mM PBS

State: Purified

State: Liquid purified IgG fraction from Bioreactor Concentrate

Stabilizer: 0.05% BSA

Preservative: 0.05% Sodium Azide

Concentration: lot specific

Purification: Protein A/G Chromatography

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 180, 145 and 125 kDa

Gene Name: neural cell adhesion molecule 1

Database Link: Entrez Gene 4684 Human


P13591

Synonyms: NCAM-1, N-CAM-1, NCAM

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Transmembrane

Protein Pathways: Cell adhesion molecules (CAMs), Prion diseases

Product images:



Formalin-paraffin human colon ganglion stained with CD56 Antibody (Clone 123A8).