

Product datasheet for **AP21442HR-N**

Chicken IgA (Fc specific) Goat Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Chicken IgA (Fc specific) Goat Polyclonal Antibody
Applications:	ELISA, ID, IF, IHC, IP
Recommended Dilution:	<p>In enzyme-immunocytochemical and immunohistochemical staining for the detection of IgA at the cellular and subcellular level by staining of appropriately treated cell and tissue substrates; to demonstrate circulating IgA antibodies in serodiagnostic microbiology and autoimmune diseases; to identify a specific antigen using a reference antibody of chicken origin known to be of the IgA isotype in the middle layer of the indirect test procedure; in non-isotopic assay methodology (e.g. ELISA) to measure IgA in chicken serum or other body fluids. Antisera to IgA do not discriminate between serum IgA (monomeric and dimeric) and higher molecular forms such as secretory IgA.</p> <p>This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.</p> <p><u>Recommended Dilutions:</u></p> <p>ELISA and comparable non-precipitating antibody-binding assays: 1/500-1/5,000. Immunocytochemistry: 1/50-1/250. Immunohistochemistry: 1/50-1/250.</p>
Reactivity:	Chicken
Host:	Goat
Immunogen:	<p>Purified normal IgA isolated from pooled Chicken serum.</p> <p>Freund's complete adjuvant is used in the first step of the immunization procedure.</p>
Isotype:	IgG
Formulation:	<p>PBS, pH 7.2 without preservatives.</p> <p>Label: HRP</p> <p>State: Lyophilized purified IgG fraction.</p> <p>Label: Horseradish Peroxidase</p> <p>Molar ratio: Peroxidase/IgG ~1.7</p>
Reconstitution Method:	Restore by adding 1 ml of sterile distilled water.



[View online »](#)

Concentration:	10 mg/ml
Purification:	Hyperimmune antisera with strong precipitating activity are selected for Fractionation by Salt-Precipitation and DEAE-Chromatography.
Conjugation:	HRP
Storage:	Store lyophilized at 2-8°C and reconstituted at 2-8°C for one week or (in aliquots) at -20°C for longer. Avoid Repeated thawing and freezing.
Note:	Adsorption: Immunoaffinity adsorbed using insolubilized antigens as required to eliminate antibodies crossreacting with other components of the immunoglobulin system or reacting with other serum proteins. Special attention is given to the removal of antibodies to common Ig/Fab. The use of insolubilized adsorption antigens prevents the presence of excess adsorbent protein or immune complexes in the antiserum.