

## **Product datasheet for AR05233PU-S**

## Interleukin-1 beta / IL-1B Chicken Protein

## **Product data:**

**Product Type:** Recombinant Proteins

**Description:** Interleukin-1 beta / IL-1B chicken recombinant protein, 50 μg

Species: Chicken

Concentration: lot specific

**Purity:** >85% by Ni chelate Chromatography

**Buffer:** State: Liquid purified protein

Buffer System: Phosphate buffered saline

**Preparation:** Liquid purified protein

**Applications:** In vitro Assay.

Functional Assays.

**Protein Description:** His-tagged recombinant chicken interleukin-1 beta expressed in *E.coli*. This antigen induces

corticosterone secretion in vivo(1) and is the immunogen used to produce chicken IL-1beta

antibody product code SP2127A.

**Storage:** Store the protein at -20°C.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: 3 month from despatch.

**RefSeq:** NP 000567

**Locus ID:** 3553

 UniProt ID:
 P01584

 Cytogenetics:
 2q14.1

Synonyms: IL-1; IL1-BETA; IL1beta; IL1F2



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



**Summary:** 

The protein encoded by this gene is a member of the interleukin 1 cytokine family. This cytokine is produced by activated macrophages as a proprotein, which is proteolytically processed to its active form by caspase 1 (CASP1/ICE). This cytokine is an important mediator of the inflammatory response, and is involved in a variety of cellular activities, including cell proliferation, differentiation, and apoptosis. The induction of cyclooxygenase-2 (PTGS2/COX2) by this cytokine in the central nervous system (CNS) is found to contribute to inflammatory pain hypersensitivity. Similarly, IL-1B has been implicated in human osteoarthritis pathogenesis. Patients with severe Coronavirus Disease 2019 (COVID-19) present elevated levels of pro-inflammatory cytokines such as IL-1B in bronchial alveolar lavage fluid samples. The lung damage induced by the Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is to a large extent, a result of the inflammatory response promoted by cytokines such as IL-1B. This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2. [provided by RefSeq, Jul 2020]

**Protein Families:** 

Druggable Genome, Secreted Protein

**Protein Pathways:** 

Alzheimer's disease, Apoptosis, Cytokine-cytokine receptor interaction, Cytosolic DNA-sensing pathway, Graft-versus-host disease, Hematopoietic cell lineage, MAPK signaling pathway, NOD-like receptor signaling pathway, Prion diseases, Toll-like receptor signaling pathway,

Type I diabetes mellitus