

## Product datasheet for **AR09784PU-L**

### Thioredoxin reductase 1 / TXNRD1 (161-649, His-tag) Human Protein

#### Product data:

<b>Product Type:</b>	Recombinant Proteins
<b>Description:</b>	Thioredoxin reductase 1 / TXNRD1 (161-649, His-tag) human recombinant protein, 0.25 mg
<b>Species:</b>	Human
<b>Expression Host:</b>	E. coli
<b>Expression cDNA Clone or AA Sequence:</b>	<u>MGSSHHHHHH</u> <u>SSGLVPRGSH</u> <u>M</u> YDYDLIIG GGSGGLAAAK EAAQYGKKVM VLDFVTPTPL GTRWGLGGTC VNVGCIPKKL MHQAALLGQA LQDSRNYGWK VEETVKHDWD RMIEAVQNHIGSLNHWGYRVA LREKKVYEN AYGQFIGPHR IKATNNKGKE KIYSAERFLI ATGERPRYLIPGDKEYCIS SDDLFLSLPYC PGKTLVVGAS YVALECAGFL AGIGLDVTVM VRSILLRFGD QDMANKIGEHEEHEGKIFIR QFVPIKVEQI EAGTPGRLRV VAQSTNSEEI IEGEYNTVML AIGRDACTRK IGLETVGVKINEKTGKIPVT DEEQTNVPIY YAIGDILEDK VELTPVAIQA GRLLAQRLYA GSTVKCDYEN VPTTVFTPLE YGACGLSEEK AVEKFGEENI EVYHSYFWPL EWTIPSRDNN KCYAKIICNT KDNERVVGFH VLGPNAGEVT QGFAAALKCG LTKKQLDSTI GIHPVCAEVF TTLSVTKRSG ASILQAGCCG
<b>Tag:</b>	His-tag
<b>Predicted MW:</b>	55.9 kDa
<b>Concentration:</b>	lot specific
<b>Purity:</b>	>90%
<b>Buffer:</b>	Presentation State: Purified State: Liquid purified protein Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 30% glycerol, 0.1 mM PMSF, 0.1M NaCl
<b>Bioactivity:</b>	Biological: Specific activity is > 15 units/mg, and was measured in a coupled assay with 5,5 - Dithiobis (2-nitrobenzoic acid) ( DTNB ) and NADPH. The amount of TNB generated by NADPH was measured in absorbance at 412 nm.
<b>Preparation:</b>	Liquid purified protein
<b>Protein Description:</b>	Recombinant human TXNRD1 protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.
<b>Storage:</b>	Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C or -70°C for longer. Avoid repeated freezing and thawing.
<b>Stability:</b>	Shelf life: one year from despatch.



[View online »](#)

RefSeq: [NP\\_001087240](#)

Locus ID: 7296

UniProt ID: [Q16881](#)

Cytogenetics: 12q23.3

Synonyms: GRIM-12; TR; TR1; TRXR1; TXNR

**Summary:** The protein encoded by this gene belongs to the pyridine nucleotide-disulfide oxidoreductase family, and is a member of the thioredoxin (Trx) system. Three thioredoxin reductase (TrxR) isozymes are found in mammals. TrxRs are selenocysteine-containing flavoenzymes, which reduce thioredoxins, as well as other substrates, and play a key role in redox homeostasis. This gene encodes an ubiquitously expressed, cytosolic form of TrxR, which functions as a homodimer containing FAD, and selenocysteine (Sec) at the active site. Sec is encoded by UGA codon that normally signals translation termination. The 3' UTRs of selenoprotein mRNAs contain a conserved stem-loop structure, the Sec insertion sequence (SECIS) element, which is necessary for the recognition of UGA as a Sec codon rather than as a stop signal. Alternative splicing, primarily at the 5' end, results in transcript variants encoding same or different isoforms, including a glutaredoxin-containing isoform that is predominantly expressed in testis. [provided by RefSeq, May 2017]

**Protein Families:** Druggable Genome

**Protein Pathways:** Pyrimidine metabolism

### Product images:

