

Product datasheet for AR50212PU-S

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

FGF10 (38-208, His-tag) Human Protein

Product data:

Product Type: Recombinant Proteins

Description: FGF10 (38-208, His-tag) human recombinant protein, 50 μg

Species: Human
Expression Host: E. coli

Expression cDNA Clone MGSSHHHHHH SSGLVPRGSH MGSHMQALGQ DMVSPEATNS SSSSFSSPSS AGRHVRSYNH

or AA Sequence: LQGDVRWRKL FSFTKYFLKI EKNGKVSGTK KENCPYSILE ITSVEIGVVA VKAINSNYYL AMNKKGKLYG

SKEFNNDCKL KERIEENGYN TYASFNWQHN GRQMYVALNG KGAPRRGQKT RRKNTSAHFL

PMVVHS

Tag: His-tag
Predicted MW: 22.0 kDa
Concentration: lot specific

Purity: >95% by SDS - PAGE

Buffer: Presentation State: Purified

State: Liquid purified protein

Buffer System: 20 mM Tris-HCl buffer (pH 8.0) containing 50% glycerol, 0.2M NaCl, 2 mM DTT,

2 mM EDTA

Endotoxin: < 1 EU per 1ug of protein (determined by LAL method)

Preparation: Liquid purified protein

Protein Description: Recombinant human FGF10 protein, fused to His-tag at N-terminus, was expressed in E.coli

and purified by using conventional chromatography.

Storage: Store undiluted at 2-8°C for one week or (in aliquots) at -20°C to -80°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

RefSeq: <u>NP 004456</u>

 Locus ID:
 2255

 UniProt ID:
 015520

 Cytogenetics:
 5p12





Summary:

The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. This protein exhibits mitogenic activity for keratinizing epidermal cells, but essentially no activity for fibroblasts, which is similar to the biological activity of FGF7. Studies of the mouse homolog of suggested that this gene is required for embryonic epidermal morphogenesis including brain development, lung morphogenesis, and initiation of lim bud formation. This gene is also implicated to be a primary factor in the process of wound healing. [provided by RefSeq, Jul 2008]


Protein Families:

Adult stem cells, Druggable Genome, Embryonic stem cells, ES Cell Differentiation/IPS, Secreted Protein, Transcription Factors, Transmembrane

Protein Pathways:

MAPK signaling pathway, Melanoma, Pathways in cancer, Regulation of actin cytoskeleton

Product images:



15% SDS-PAGE (3ug)