

Product datasheet for **CF500516**

Grp75 (HSPA9) Mouse Monoclonal Antibody [Clone ID: OTI9F8]

Product data:

| | |
|-------------------------|--|
| Product Type: | Primary Antibodies |
| Clone Name: | OTI9F8 |
| Applications: | FC, IF, IHC, WB |
| Recommended Dilution: | WB 1:1000~2000, IHC 1:50, IF 1:50, FLOW 1:100 |
| Reactivity: | Human, Dog, Rat, Monkey, Mouse |
| Host: | Mouse |
| Isotype: | IgG2b |
| Clonality: | Monoclonal |
| Immunogen: | Full length human recombinant protein of human HSPA9 (NP_004125) produced in HEK293T cell. |
| Formulation: | Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose) |
| Reconstitution Method: | For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific) |
| Purification: | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G) |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C as received. |
| Stability: | Stable for 12 months from date of receipt. |
| Predicted Protein Size: | 73.5 kDa |
| Gene Name: | heat shock protein family A (Hsp70) member 9 |
| Database Link: | NP_004125 Entrez Gene 15526 Mouse Entrez Gene 291671 Rat Entrez Gene 474697 Dog Entrez Gene 714506 Monkey Entrez Gene 3313 Human P38646 |



[View online »](#)

Background:

This gene encodes a member of the heat shock protein 70 gene family. The encoded protein is primarily localized to the mitochondria but is also found in the endoplasmic reticulum, plasma membrane and cytoplasmic vesicles. This protein is a heat-shock cognate protein. This protein plays a role in cell proliferation, stress response and maintenance of the mitochondria. A pseudogene of this gene is found on chromosome 2.

Synonyms:


CSA; GRP-75; GRP75; HEL-S-124m; HSPA9B; MOT; MOT2; MTHSP75; PBP74

Protein Families:


Stem cell - Pluripotency

Protein Pathways:


RNA degradation

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY HSPA9 (Cat# [RC201397], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HSPA9 (Cat# [TA500516]). Positive lysates [LY401334] (100ug) and [LC401334] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-HSPA9 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).




Western blot analysis of extracts (10ug) from a mouse cell line and 3 different mouse tissues by using anti-HSPA9 monoclonal antibody (1:200).




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human breast tissue using anti-HSPA9 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500516])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human kidney tissue using anti-HSPA9 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500516])




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-HSPA9 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500516])




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-HSPA9 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500516])




Immunohistochemical staining of paraffin-embedded Carcinoma of Human pancreas tissue using anti-HSPA9 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500516])




Immunohistochemical staining of paraffin-embedded Human endometrium tissue within the normal limits using anti-HSPA9 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA500516])




Anti-HSPA9 mouse monoclonal antibody ([TA500516]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY HSPA9 ([RC201397]).



Immunofluorescent staining of HepG2 cells using anti-HSPA9 mouse monoclonal antibody ([TA500516]).



Flow cytometric Analysis of HeLa cells, using anti-HSPA9 antibody ([TA500516]), (Red), compared to a nonspecific negative control antibody, (Blue).



Flow cytometric Analysis of Jurkat cells, using anti-HSPA9 antibody ([TA500516]), (Red), compared to a nonspecific negative control antibody, (Blue).