

Product datasheet for CF502407

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

beta Catenin (CTNNB1) Mouse Monoclonal Antibody [Clone ID: OTI3F9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3F9

Applications: FC, IHC, WB

Recommended Dilution: WB 1:500~2000, IHC 1:150, FLOW 1:100

Reactivity: Human, Dog, Rat, Monkey, Mouse

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CTNNB1 (NP_001895) produced in

HEK293T cell.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 85.3 kDa

Gene Name: catenin beta 1

Database Link: NP 001895

Entrez Gene 12387 MouseEntrez Gene 84353 RatEntrez Gene 477032 DogEntrez Gene 574265

MonkeyEntrez Gene 1499 Human

P35222





Background:

The protein encoded by this gene is part of a complex of proteins that constitute adherens junctions (AJs). AJs are necessary for the creation and maintenance of epithelial cell layers by regulating cell growth and adhesion between cells. The encoded protein also anchors the actin cytoskeleton and may be responsible for transmitting the contact inhibition signal that causes cells to stop dividing once the epithelial sheet is complete. Finally, this protein binds to the product of the APC gene, which is mutated in adenomatous polyposis of the colon. Mutations in this gene are a cause of colorectal cancer (CRC), pilomatrixoma (PTR), medulloblastoma (MDB), and ovarian cancer. Three transcript variants encoding the same protein have been found for this gene.

Synonyms: armadillo; CTNNB; MRD19


Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Transcription Factors


Protein Pathways: Adherens junction, Arrhythmogenic right ventricular cardiomyopathy (ARVC), Basal cell carcinoma, Colorectal cancer, Endometrial cancer, Focal adhesion, Leukocyte transendothelial


migration, Melanogenesis, Pathogenic Escherichia coli infection, Pathways in cancer, Prostate

cancer, Thyroid cancer, Tight junction, Wnt signaling pathway


Product images:









Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-CTNNB1 monoclonal antibody.



Flow cytometric Analysis of Hela cells, using anti-CTNNB1 antibody ([TA502407]), (Red), compared to a nonspecific negative control antibody, (Blue).




HEK293T cells transfected with either [RC208947] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-CTNNB1 antibody ([TA502407]), and then analyzed by flow cytometry.




Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-CTNNB1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502407])






Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-CTNNB1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502407])



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-CTNNB1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, [TA502407])



Western blot analysis of extracts (10ug) from 9 Human tissue by using anti-CTNNB1 monoclonal antibody at 1:200 (1: Testis; 2: Omentum; 3: Uterus; 4: Breast; 5: Brain; 6: Liver; 7: Ovary; 8: Thyroid gland; 9: Colon).