

## **Product datasheet for CF802546**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### Ki67 (MKI67) Mouse Monoclonal Antibody [Clone ID: OTI7G8]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI7G8

Applications: IHC

Recommended Dilution: IHC 1:100

Reactivity: Human Host: Mouse

Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 1160-1493 of human

MKI67 (NP\_002408) produced in E.coli.

**Formulation:** Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

**Reconstitution Method:** For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** marker of proliferation Ki-67

Database Link: NP 002408

Entrez Gene 4288 Human

P46013

**Background:** This gene encodes a nuclear protein that is associated with and may be necessary for cellular

proliferation. Alternatively spliced transcript variants have been described. A related

pseudogene exists on chromosome X. [provided by RefSeq, Mar 2009]

Synonyms: KIA; MIB-; MIB-1; PPP1R105






**Protein Families:** 

Druggable Genome, ES Cell Differentiation/IPS

# **Product images:**



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-MKI67 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA802546])