

Product datasheet for CF802721

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PARN Mouse Monoclonal Antibody [Clone ID: OTI2D9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2D9
Applications: IHC, WB
Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-266 of human PARN

(NP 002573) produced in E.coli.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 73.3 kDa

Gene Name: poly(A)-specific ribonuclease

Database Link: NP 002573

Entrez Gene 74108 MouseEntrez Gene 5073 Human

<u>095453</u>





Background:


The protein encoded by this gene is a 3'-exoribonuclease, with similarity to the RNase D family of 3'-exonucleases. It prefers poly(A) as the substrate, hence, efficiently degrades poly(A) tails of mRNAs. Exonucleolytic degradation of the poly(A) tail is often the first step in the decay of eukaryotic mRNAs. This protein is also involved in silencing of certain maternal mRNAs during oocyte maturation and early embryonic development, as well as in nonsensemediated decay (NMD) of mRNAs that contain premature stop codons. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Aug 2008]

Synonyms: DAN


Protein Families: Transcription Factors

Protein Pathways: RNA degradation

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PARN ([RC207220], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PARN. Positive lysates [LY419233] (100ug) and [LC419233] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-PARN mouse monoclonal antibody. ([TA802721]) Dilution: 1:150





Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-PARN mouse monoclonal antibody. ([TA802721]) Dilution: 1:150