

Product datasheet for **CF806835**

PD1 (PDCD1) Mouse Monoclonal Antibody [Clone ID: OTI8B1]

Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI8B1
Applications:	ELISA, FC, IF, LMNX, Neutralize, WB
Recommended Dilution:	WB 1:2000, IF 1:100~900, FLOW 1:50
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human PDCD1 (NP_005009) produced in HEK293T cell.
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	29.2 kDa
Gene Name:	programmed cell death 1
Database Link:	NP_005009 Entrez Gene 5133 Human Q15116
Synonyms:	CD279; hPD-1; hPD-I; hSLE1; PD-1; PD1; SLEB2
Protein Families:	Druggable Genome, Transmembrane




[View online »](#)

Protein Pathways: Cell adhesion molecules (CAMs), T cell receptor signaling pathway

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PDCD1 ([RC210364], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PDCD1. Positive lysates [LY401555] (100ug) and [LC401555] (20ug) can be purchased separately from OriGene.



PD1 ELISA with 1B11 Capture ([CF807867]) and 8B1 Detection (CF806835) Antibodies. Substrate used: Recombinant Human PD1 ([TP310364])

PD1 cell line


293T cell line




Immunofluorescent staining of PDCD1 (RC210364)-stable-transfected HEK293T cells (left) labeling PDCD1 with mouse monoclonal antibody [TA806835] (green) and nucleus with Hoechst33342 (blue). HEK293T cells serve as negative control (right). n (1:100)

PD1 cell line


293T cell line




Immunocytochemistry staining of stable expression PD1 cells using anti-PDCD1 mouse monoclonal antibody ([TA806835]) (Left). The right is negative control. (1:100) (1:900)




Flow cytometric Analysis of stable expression PD1 cells using anti-PDCD1 antibody ([TA806835]) (Red) compared to a nonspecific negative control antibody (Blue) (1:50).




HEK293T cells transfected with either mouse PD1 ([MR227347]) overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-PDCD1 antibody ([TA806835]), compared to a nonspecific negative control antibody (green), and then analyzed by flow cytometry (1:50).



PD1 Luminex ELISA with 1B11 Capture ([CF807867]) and 8B1 Detection (CF806835) Antibodies. Substrate used: Recombinant Human PD1 ([TP310364])



PD1 Luminex ELISA with 7B4 Capture ([CF807995]) and 8B1 Detection (CF806835) Antibodies. Substrate used: Recombinant Human PD1 ([TP310364])



Flow cytometric Analysis of stable expression PDL1 ([RC213071]) cells using anti-PDCD1 antibody ([TA806835]) (blue) or 0.3ug/ml PD1-Fc fusion protein ([TP700199]) (green) or both (red), and detected by anti-Fc (human) IgG-FITC (1:50).