

Product datasheet for CF809362

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

n-Myc (MYCN) Mouse Monoclonal Antibody [Clone ID: OTI1D6]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1D6

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 229-464 of human

MYCN(NP 005369) produced in E.coli.

Formulation: Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)

Reconstitution Method: For reconstitution, we recommend adding 100uL distilled water to a final antibody

concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 49.4 kDa

Gene Name: MYCN proto-oncogene, bHLH transcription factor

Database Link: NP 005369

Entrez Gene 4613 Human

P04198





Background: This gene is a member of the MYC family and encodes a protein with a basic helix-loop-helix


(bHLH) domain. This protein is located in the nucleus and must dimerize with another bHLH protein in order to bind DNA. Amplification of this gene is associated with a variety of tumors, most notably neuroblastomas. Multiple alternatively spliced transcript variants encoding

different isoforms have been found for this gene. [provided by RefSeq, Jun 2014]

Synonyms: bHLHe37; MODED; N-myc; NMYC; ODED

Protein Families: Druggable Genome, Transcription Factors

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MYCN ([RC201241], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MYCN (1:2000). Positive lysates [LY417334] (100ug) and [LC417334] (20ug) can be purchased separately from OriGene.