

## Product datasheet for **CF813206**

### CD70 Mouse Monoclonal Antibody [Clone ID: OTI5E9]

#### Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI5E9
Applications:	FC
Recommended Dilution:	FLOW 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human CD70 (NP_001243) produced in HEK293T cell.
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	20.9 kDa
Gene Name:	CD70 molecule
Database Link:	<a href="#">NP_001243</a> <a href="#">Entrez Gene 970 Human</a> <a href="#">P32970</a>
Background:	Cytokine that binds to CD27. Plays a role in T-cell activation. Induces the proliferation of costimulated T-cells and enhances the generation of cytolytic T-cells. [UniProtKB/Swiss-Prot Function]




[View online »](#)

**Synonyms:** CD27-L; CD27L; CD27LG; LPFS3; TNFSF7; TNLG8A  
**Protein Families:** ES Cell Differentiation/IPS, Transmembrane  
**Protein Pathways:** Cytokine-cytokine receptor interaction


**Product images:**





Flow cytometric analysis of living 293T cells transfected with CD70 overexpression plasmid ([RC200410]), Red/empty vector ([PS100001], Blue) using anti-CD70 antibody ([TA813206]). Cells incubated with a non-specific antibody (Green) were used as isotype control (1:100).





Flow cytometric analysis of living Raji cells, using anti-CD70 antibody ([TA813206], Red), compared to an isotype control (green), and a PBS control (blue) (1:100).



Flow cytometric analysis of living 786-O cells, using anti-CD70 antibody ([TA813206], Red), compared to an isotype control (green), and a PBS control (blue) (1:100).



Flow cytometric analysis of living PBMCs treated with 10ug/ml PHA for 72h (Right)/untreated (Left) using anti-CD70 antibody ([TA813206]) (1:100).



Flow cytometric analysis of living PBMCs treated with 10ug/ml PHA for 72h using anti-CD70 antibody ([TA813206]) (Right). Cells incubated with a non-specific antibody (Left) were used as isotype control (1:100).