

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for DDX9022A488-100

CD49b (ITGA2) Human Monoclonal Antibody [Clone ID: B4F12]

Product data:

Product Type: Primary Antibodies

Clone Name: B4F12
Applications: FC, IF

Recommended Dilution: IF, ELISA, preferential detection of activated platelet by flow cytometry, platelet

aggregation studies.

Usage recommendation:

*This monoclonal antibody may be used between 5-25µg/ml.

*Optimal dilution should be determined by each laboratory for each application.

*Coupled antibody: to maintain RT before use.

Reactivity: Human
Host: Human
Isotype: IgG1

Clonality: Monoclonal

Immunogen: CD40 activated and EBV immortalized human B lymphocytes

Specificity: GPIa-IIa preferentially on acitvated platelets.

Formulation: 100 μ g in 200 μ l / 50 μ g in 100 μ l PBS 50% glycerol

Label: Alexa Fluor 488

Concentration: lot specific

Purification: QMA Hyper D ion exchange chromatography

Conjugation: Alexa Fluor 488

Storage: -20°C. KEEP CONTENTS STERILE: no preservative.

Purified antibodies: avoid repeated freeze/thaw cycles. Coupled antibodies: glycerol protects from freezing.

Gene Name: integrin subunit alpha 2

Database Link: <u>Entrez Gene 3673 Human</u>






Background:

Neonatal AlloImmune Thrombocytopenia (NAIT) is the commonest cause of severe isolated thrombocytopenia in the foetus and newborn. In Caucasians, 97% of NAIT are caused by HPA-1a or HPA-5b alloantibodies. B4F12 was isolated after combination of CD40 activation and Epstein-Barr virus transformation of B cells isolated from a woman with high titre of anti - HPA-5b alloantibodies. B4F12 human monoclonal antibody recognizes GPIa-IIa preferentially on activated human platelets (Mérieux Y, Rigal D, Pin JJ, XVIIth Regional Congress, International Society of Blood Transfusion, Madrid 2007; Guillot-Chene P, Lebecque S, Rigal D, Annales Pharmaceutiques Françaises, 06/2009, 67 (3) 182-6).

Synonyms:

Integrin alpha-2, ITGA-2, VLA-2 alpha, VLA2, GPIa, Collagen Receptor


Product images:



IF staining of human platelets with **B4F12**



ELISA detection of human platelets with **B4F12**



Inhibition of collagen-induced platelet aggregation by **F2H11** and **B4F12**