

Product datasheet for KN312748LP

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US

OriGene Technologies, Inc.

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Padi4 Mouse Gene Knockout Kit (CRISPR)

Product data:

Product Type: Knockout Kits (CRISPR)

Format: 2 gRNA vectors, 1 Luciferase-Puro donor, 1 scramble control

Donor DNA: Luciferase-Puro

Symbol: Padi4 Locus ID: 18602

Components: KN312748G1, Padi4 gRNA vector 1 in pCas-Guide CRISPR vector (GE100002)

KN312748G2, Padi4 gRNA vector 2 in pCas-Guide CRISPR vector (GE100002)

KN312748LPD, donor DNA containing left and right homologous arms and Luciferase-Puro

functional cassette.

GE100003, scramble sequence in pCas-Guide vector

Disclaimer: These products are manufactured and supplied by OriGene under license from ERS. The kit is

designed based on the best knowledge of CRISPR technology. The system has been functionally validated for knocking-in the cassette downstream the native promoter. The efficiency of the knock-out varies due to the nature of the biology and the complexity of the

experimental process.

 RefSeq:
 NM 011061

 UniProt ID:
 Q9Z183

Synonyms: Pad4; Pdi4

Summary: Catalyzes the citrullination/deimination of arginine residues of proteins such as histones,

thereby playing a key role in histone code and regulation of stem cell maintenance. Citrullinates histone H1 at 'Arg-54' (to form H1R54ci), histone H3 at 'Arg-2', 'Arg-8', 'Arg-17' and/or 'Arg-26' (to form H3R2ci, H3R8ci, H3R17ci, H3R26ci, respectively) and histone H4 at 'Arg-3' (to form H4R3ci). Acts as a key regulator of stem cell maintenance by mediating citrullination of histone H1: citrullination of 'Arg-54' of histone H1 (H1R54ci) results in H1 displacement from chromatin and global chromatin decondensation, thereby promoting pluripotency and stem cell maintenance. Promotes profound chromatin decondensation during the innate immune response to infection in neutrophils by mediating formation of

H1R54ci. Citrullination of histone H3 prevents their methylation by CARM1 and

HRMT1L2/PRMT1 and represses transcription. Citrullinates EP300/P300 at 'Arg-2142', which

favors its interaction with NCOA2/GRIP1.[UniProtKB/Swiss-Prot Function]





Product images:

Donor Vector Edited Chromosome



RFP, Luc, and mBFP will be under native gene promoter