

Product datasheet for LC400973

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SAPK4 (MAPK13) (NM 002754) Human Over-expression Lysate

Product data:

Product Type: Over-expression Lysates

Description: MAPK13 HEK293T cell transient overexpression lysate (as WB positive control)

Species: Human HEK293T **Expression Host:**

Expression cDNA Clone

or AA Sequence:

TrueORF Clone RC200606

Tag: C-Myc/DDK

Detection Antibodies: Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)

ACCN: NM 002754, NP 002745

MAPK-13; MAPK 13; p38delta; PRKM13; SAPK4 Synonyms:

Predicted MW: 42.1 kDa

Components: 1 vial of 20 ug lyophilized gene specific transient over-expression cell lysate

The lysate can be shipped at ambient temperature. Upon receiving, store the sample at -Storage:

> 20°C. Lysate samples can be reconstituted with SDS Sample Buffer. Avoid repeated freezethaw cycles after reconstitution. Lysate samples are stable for 12 months from date of receipt

when stored at -20°C.

Preparation: HEK293T cells in 10-cm dishes were transiently transfected with MegaTran Transfection

> Reagent (TT200002) and 5ug TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). To facilitate transportation and protein, the

products are supplied as lyophilized proteins.

NP 002745 RefSeq:

Locus ID: 5603 **Cytogenetics:** 6p21.31

Protein Families: Druggable Genome, Protein Kinase






Protein Pathways:

Amyotrophic lateral sclerosis (ALS), Epithelial cell signaling in Helicobacter pylori infection, Fc epsilon RI signaling pathway, GnRH signaling pathway, Leukocyte transendothelial migration, MAPK signaling pathway, Neurotrophin signaling pathway, NOD-like receptor signaling pathway, Progesterone-mediated oocyte maturation, RIG-I-like receptor signaling pathway, T cell receptor signaling pathway, Toll-like receptor signaling pathway

Product images:



Western blot validation of overexpression lysate (Cat# [LY400973]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC200606] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).