

## Product datasheet for LC429361

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## beta TRCP2 (FBXW11) (NM\_012300) Human Over-expression Lysate

**Product data:** 

**Product Type:** Over-expression Lysates

**Description:** FBXW11 HEK293T cell transient overexpression lysate (as WB positive control)

Species: Human
Expression Host: HEK293T

**Expression cDNA Clone** 

TrueORF Clone RC218905

or AA Sequence:

Tag: C-Myc/DDK

**Detection Antibodies:** Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)

ACCN: <u>NM 012300, NP 036432</u>

Synonyms: BTRC2; BTRCP2; FBW1B; Fbw11; FBXW1B; Hos; NEDJED

**Predicted MW:** 61.9 kDa

**Components:** 1 vial of 20 ug lyophilized gene specific transient over-expression cell lysate

Storage: The lysate can be shipped at ambient temperature. Upon receiving, store the sample at -

20°C. Lysate samples can be reconstituted with SDS Sample Buffer. Avoid repeated freezethaw cycles after reconstitution. Lysate samples are stable for 12 months from date of receipt

when stored at -20°C.

**Preparation:** HEK293T cells in 10-cm dishes were transiently transfected with MegaTran Transfection

Reagent (TT200002) and 5ug <u>TrueORF</u> cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). To facilitate transportation and protein, the

products are supplied as lyophilized proteins.

**RefSeq:** NP 036432

**Locus ID:** 23291 **Cytogenetics:** 5q35.1

**Protein Families:** Druggable Genome





## beta TRCP2 (FBXW11) (NM\_012300) Human Over-expression Lysate - LC429361

**Protein Pathways:** 

Hedgehog signaling pathway, Oocyte meiosis, Ubiquitin mediated proteolysis, Wnt signaling pathway