

Product datasheet for **LY402895**

NF-kB p65 (RELA) (NM_021975) Human Over-expression Lysate

Product data:



Product Type:	Over-expression Lysates
Description:	Transient overexpression lysate of v-rel reticuloendotheliosis viral oncogene homolog A (avian) (RELA), transcript variant 1
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	TrueORF Clone RC220780
Tag:	C-Myc/DDK
Detection Antibodies:	Clone OTI4C5, Anti-DDK (FLAG) monoclonal antibody (TA50011-100)
ACCN:	NM_021975 , NP_068810
Synonyms:	CMCU; NFKB3; p65
Predicted MW:	60 kDa
Components:	1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT)
Storage:	The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at -80°C.
Bioactivity:	RELA Activity Verified in a DNA-binding Assay
Preparation:	HEK293T cells in 10-cm dishes were transiently transfected with MegaTran Transfection Reagent (TT200002) and 5ug TrueORE cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix (Sigma), 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit (Thermo Scientific Inc.). Cell lysates were aliquoted and stored at -20°C before shipping.



[View online »](#)

RefSeq: [NP_068810](#)
Locus ID: 5970
Cytogenetics: 11q13.1
Protein Families: Druggable Genome, Transcription Factors
Protein Pathways: Acute myeloid leukemia, Adipocytokine signaling pathway, Apoptosis, B cell receptor signaling pathway, Chemokine signaling pathway, Chronic myeloid leukemia, Cytosolic DNA-sensing pathway, Epithelial cell signaling in Helicobacter pylori infection, MAPK signaling pathway, Neurotrophin signaling pathway, NOD-like receptor signaling pathway, Pancreatic cancer, Pathways in cancer, Prostate cancer, RIG-I-like receptor signaling pathway, Small cell lung cancer, T cell receptor signaling pathway, Toll-like receptor signaling pathway

Product images:



RELA activity in the over-expression lysate (LY402895) and a vector-transfected control lysate was measured in a colorimetric DNA-binding assay. Double-stranded oligonucleotide containing the RELA consensus DNA-binding sequence was incubated with dilutions of the over-expression lysate and RELA bound to the oligo was captured onto the surface of a microtiter plate. After washing, bound RELA was detected with an anti-RELA primary antibody followed by an HRP-labeled secondary antibody. After initial color development, the reaction was quenched and the color intensity was measured at 450nm. The data show high levels of RELA DNA binding activity in the over-expression lysate, but almost no DNA-binding activity in the control lysate. Overexpression cell lysates are prepared from HEK293T cells transfected with [RC220780] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

Western blot validation of overexpression lysate (Cat# LY402895) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC220780] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).