

Product datasheet for **MC204684**

Anapc13 (NM_181394) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Anapc13 (NM_181394) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Anapc13
Synonyms:	1810004D07Rik; APC13; SWM1
Mammalian Cell Selection:	Neomycin
Vector:	PCMV6-Kan/Neo (PCMV6KN)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>BC028526 GACGCGCGCAGTGGGCTGACAAGATCAAAGCTGCAGGAGGATGGACAGTGAGGTACAGCGAGATGGA AGGATCTTGGACCTGATTGATGATGCTTGGCGGAAGATAAGCTGCCATATGAGGATGTCGCCATCCAC TGAGTGAGCTTCTGAGCCCGAGCAAGACAACGGAGGCACCACAGAGTCTGTGAAAGAACAGGAGATGAA GTGGACAGACCTGGCCTTACAGGGCCTCCACGAGAACGTCCCACCGCTGAAAAGTATGCCGGGCTCCT TCCCACGGACGGAGTTTTCAAACATATGGGGATAAAAAATGTTTCTGTGCCAAATGCAGATCTTTAGCA ATAAATCAACACTCAAAAGCATAAAAAAAAAAAAAAAAAAAAAA
Restriction Sites:	EcoRI-NotI
ACCN:	NM_181394
Insert Size:	225 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



[View online »](#)

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [BC028526](#), [AAH28526](#)

RefSeq Size: 395 bp

RefSeq ORF: 225 bp

Locus ID: 69010

UniProt ID: [Q8R034](#)

Cytogenetics: 9 F1

Gene Summary: Component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle. The APC/C complex acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains (By similarity). [UniProtKB/Swiss-Prot Function]