

# Product datasheet for MG226484

## Ddx3x (NM\_010028) Mouse Tagged ORF Clone

### **Product data:**

Product Type:	Expression Plasmids
Product Name:	Ddx3x (NM_010028) Mouse Tagged ORF Clone
Tag:	TurboGFP
Symbol:	Ddx3x
Synonyms:	D1Pas1-rs2; Ddx3; Fin14
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-AC-GFP (PS100010)
E. coli Selection:	Ampicillin (100 ug/mL)

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

DRF Nucleotide Sequence:	>MG226484 representing NM_010028 Red=Cloning site Blue=ORF Green=Tags(s)
	TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C
	ATGAGTCATGTGGCAGTGGAAAATGCGCTCGGGCTGGACCAGCAGTTTGCTGGCCTAGACCTGAACTCTT
	CAGATAATCAGAGTGGAGGAAGTACAGCAAGCAAAGGGCGTTATATCCCACCTCATTTAAGGAACAGAGA
	AGCTACTAAAGGATTCTATGACAAAGACAGTTCAGGGTGGAGTTCTAGTAAAGATAAGGATGCATACAGC
	AGTTTTGGATCACGGGGTGATTCAAGAGGGAAGTCTAGCTTCTTTGGAGATCGTGGAAGTGGATCAAGGG
	GAAGGTTTGATGATCGTGGACGGGGAGACTATGATGGCATTGGTGGCCGTGGAGATAGAAGTGGCTTTGG
	CAAATTTGAAAGAGGTGGAAATAGTCGCTGGTGTGACAAATCAGATGAAGATGACTGGTCAAAGCCACTC
	CCACCAAGTGAACGATTGGAACAGGAACTCTTTTCTGGAGGCAATACTGGGATTAACTTTGAGAAATATG
	ATGACATTCCAGTCGAAGCAACAGGCAACAACTGTCCTCCACACATTGAAAGTTTCAGTGATGTCGAGAT
	GGGAGAAATTATTATGGGAAACATTGAGCTTACTCGTTATACTCGCCCAACTCCAGTGCAGAAGCATGCT
	ATTCCTATTATCAAAGAGAAAAGAGACTTGATGGCTTGTGCTCAAACAGGCTCTGGAAAAACTGCAGCAT
	TTCTCTTGCCCATCTTGAGTCAGATCTATGCTGATGGTCCAGGAGAAGCTCTGAGGGCTATGAAGGAAAA
	TGGAAGATATGGCCGTCGTAAACAGTATCCAATCTCTTTGGTACTGGCACCAACGAGAGAATTGGCAGTG
	CAGATCTATGAGGAAGCCAGAAAATTCTCATACCGATCTAGAGTCCGTCC
	CTGAAATTGGCCAGCAGATTCGAGACTTAGAACGTGGATGCCACTTGTTAGTAGCCACTCCAGGACGTCT
	AGTGGATATGATGGAGAGAGGGAAGATCGGGTTAGACTTCTGCAAATACCTGGTGTTAGATGAAGCTGAC
	CGGATGTTAGATATGGGGTTTGAACCTCAGATACGAAGAATAGTTGAACAAGACACTATGCCTCCAAAAG
	GTGTCCGCCACACTATGATGTTTAGTGCTACTTTTCCTAAGGAAATACAGATGCTGGCCCGTGATTTCTT
	AGATGAGTACATATTTCTGGCTGTAGGAAGAGTTGGGTCTACTTCAGAGAACATCACAAAAAGTGGTT
	TGGGTGGAGGAGATAGACAAAAGGTCATTTCTGCTTGACCTTCTAAATGCAACAGGCAAGGATTCCCTGA
	CTCTAGTGTTTGTGGAGACCAAAAAGGGGGGCAGATTCGCTGGAGGATTTCTTATACCATGAAGGATATGC
	TTGTACCAGTATCCATGGAGACCGTTCTCAGAGAGAGAGA
	AAAAGCCCAATTCTAGTGGCTACAGCAGTAGCAGCAAGAGGACTGGATATTTCAAATGTGAAGCATGTTA
	TTAATTTTGACCTGCCTAGTGATATCGAAGAATATGTGCATCGCATAGGCCGTACAGGCCGTGTGGGAAA
	CCTTGGTCTTGCCACCTCATTCTTTAATGAAAGGAATATAAATATCACAAAGGATTTACTGGATCTTCTT
	GTTGAAGCAAAACAAGAAGTGCCTTCTTGGTTAGAGAACATGGCTTTTGAACACCACTACAAGGGTAGCA

ACGCGTACGCGGCCGCTCGAG - GFP Tag - GTTTAA

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Ddx3x (NM\_010028) Mouse Tagged ORF Clone - MG226484

Protein Sequence:

>MG226484 representing NM\_010028
Red=Cloning site Green=Tags(s)

MSHVAVENALGLDQQFAGLDLNSSDNQSGGSTASKGRYIPPHLRNREATKGFYDKDSSGWSSSKDKDAYS SFGSRGDSRGKSSFFGDRGSGSRGRFDDRGRGDYDGIGGRGDRSGFGKFERGGNSRWCDKSDEDDWSKPL PPSERLEQELFSGGNTGINFEKYDDIPVEATGNNCPPHIESFSDVEMGEIIMGNIELTRYTRPTPVQKHA IPIIKEKRDLMACAQTGSGKTAAFLLPILSQIYADGPGEALRAMKENGRYGRRKQYPISLVLAPTRELAV QIYEEARKFSYRSRVRPCVVYGGAEIGQQIRDLERGCHLLVATPGRLVDMMERGKIGLDFCKYLVLDEAD RMLDMGFEPQIRRIVEQDTMPPKGVRHTMMFSATFPKEIQMLARDFLDEYIFLAVGRVGSTSENITQKVV WVEEIDKRSFLLDLLNATGKDSLTLVFVETKKGADSLEDFLYHEGYACTSIHGDRSQRDREEALHQFRSG KSPILVATAVAARGLDISNVKHVINFDLPSDIEEYVHRIGRTGRVGNLGLATSFFNERNINITKDLLDLL VEAKQEVPSWLENMAFEHHYKGSSRGRSKSSRFSGGFGARDYRQSSGASSSSFSSSRASSSRSGGGHGG SRGFGGGGYGGFYNSDGYGGNYNSQGVDWWGN

TRTRPLE - GFP Tag - V

Sgfl-Mlul

Restriction Sites:

### Cloning Scheme:



ACCN:	NM_010028
ORF Size:	1986 bp
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

# **Ddx3x (NM\_010028) Mouse Tagged ORF Clone – MG226484**

Reconstitution Method:	<ol> <li>Centrifuge at 5,000xg for 5min.</li> <li>Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>Close the tube and incubate for 10 minutes at room temperature.</li> <li>Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM 010028.3, NP 034158.1</u>
RefSeq Size:	4571 bp
RefSeq ORF:	1989 bp
Locus ID:	13205
UniProt ID:	<u>Q62167</u>
Cytogenetics:	X 8.17 cM

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### 

#### Ddx3x (NM\_010028) Mouse Tagged ORF Clone - MG226484

#### Gene Summary:

Multifunctional ATP-dependent RNA helicase. The ATPase activity can be stimulated by various ribo- and deoxynucleic acids indicative for a relaxed substrate specificity. In vitro can unwind partially double-stranded DNA with a preference for 5'-single-stranded DNA overhangs. Is involved in several steps of gene expression, such as transcription, mRNA maturation, mRNA export and translation. However, the exact mechanisms are not known and some functions may be specific for a subset of mRNAs. Involved in transcriptional regulation. Can enhance transcription from the CDKN1A/WAF1 promoter in a SP1-dependent manner. Found associated with the E-cadherin promoter and can down-regulate transcription from the promoter. Involved in regulation of translation initiation. Proposed to be involved in positive regulation of translation such as of cyclin E1/CCNE1 mRNA and specifically of mRNAs containing complex secondary structures in their 5'UTRs; these functions seem to require RNA helicase activity. Specifically promotes translation of a subset of viral and cellular mRNAs carrying a 5'proximal stem-loop structure in their 5'UTRs and cooperates with the eIF4F complex. Proposed to act prior to 43S ribosomal scanning and to locally destabilize these RNA structures to allow recognition of the mRNA cap or loading onto the 40S subunit. After association with 40S ribosomal subunits seems to be involved in the functional assembly of 80S ribosomes; the function seems to cover translation of mRNAs with structured and nonstructured 5'UTRs and is independent of RNA helicase activity. Also proposed to inhibit capdependent translation by competetive interaction with EIF4E which can block the EIF4E:EIF4G complex formation. Proposed to be involved in stress response and stress granule assembly; the function is independent of RNA helicase activity and seems to involve association with EIF4E. May be involved in nuclear export of specific mRNAs but not in bulk mRNA export via interactions with XPO1 and NXF1. Also associates with polyadenylated mRNAs independently of NXF1. Associates with spliced mRNAs in an exon junction complex (EJC)-dependent manner and seems not to be directly involved in splicing. May be involved in nuclear mRNA export by association with DDX5 and regulating its nuclear location. Involved in innate immune signaling promoting the production of type I interferon (IFN-alpha and IFN-beta); proposed to act as viral RNA sensor, signaling intermediate and transcriptional coactivator. Involved in TBK1 and IKBKE-dependent IRF3 activation leading to IFNB induction, plays a role of scaffolding adapter that links IKBKE and IRF3 and coordinates their activation. Also found associated with IFNB promoters; the function is independent of IRF3. Can bind to viral RNAs and via association with MAVS/IPS1 and DDX58/RIG-I is thought to induce signaling in early stages of infection. Involved in regulation of apoptosis. May be required for activation of the intrinsic but inhibit activation of the extrinsic apoptotic pathway. Acts as an antiapoptotic protein through association with GSK3A/B and BIRC2 in an apoptosis antagonizing signaling complex; activation of death receptors promotes caspase-dependent cleavage of BIRC2 and DDX3X and relieves the inhibition. May be involved in mitotic chromosome segregation. Is an allosteric activator of CSNK1E, it stimulates CSNK1E-mediated phosphorylation of DVL2 and is involved in the positive regulation of canonical Wnt signaling (By similarity).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



## **Product images:**



Circular map for MG226484

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US