

Product datasheet for MR201190L3

Cbx7 (NM_144811) Mouse Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Cbx7 (NM_144811) Mouse Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	Cbx7
Synonyms:	1600014J01Rik; AI851678; D15Ert417e
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR201190).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.


ACCN:	NM_144811
ORF Size:	477 bp



[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_144811.3 , NP_659060.1
RefSeq Size:	2893 bp
RefSeq ORF:	477 bp
Locus ID:	52609
UniProt ID:	Q8VDS3
Cytogenetics:	15 37.85 cM
Gene Summary:	Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development (PubMed:16537902, PubMed:22226355). PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility. Promotes histone H3 trimethylation at 'Lys-9' (H3K9me3) (By similarity). Binds to histone H3 trimethylated at 'Lys-9' (H3K9me3) or at 'Lys-27' (H3K27me3) (PubMed:16537902, PubMed:22226355). Trimethylation at 'Lys-27' (H3K27me3) is important for chromatin recruitment (PubMed:22226355, PubMed:16537902). May possibly also bind trimethylated lysine residues in other proteins (in vitro) (PubMed:16537902). Binds non-coding, single-stranded RNA and double-stranded RNA (PubMed:20541999, PubMed:16537902). Plays a role in the timely repression of differentiation-specific genes in pluripotent embryonic stem cells to maintain the undifferentiated state (PubMed:22226355). Regulator of cellular lifespan by maintaining the repression of CDKN2A, but not by inducing telomerase activity (PubMed:14647293).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR201190L3