

# Product datasheet for MR208047L4

# Traf2 (NM\_009422) Mouse Tagged Lenti ORF Clone

# **Product data:**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Expression Plasmids
Product Name:	Traf2 (NM_009422) Mouse Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	Traf2
Synonyms:	AI325259
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR208047).
<b>Restriction Sites:</b>	Sgfl-Mlul
Cloning Scheme:	
	Cloning sites used for ORF Shuttling:
	Sgf I         ORF         Mlu I            GCG ATC GC         ATG NNN         ACG CGT



ACCN: ORF Size: NM\_009422 1503 bp



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


<b>CRIGENE</b> Traf2 (NM_009422) Mouse Tagged Lenti ORF Clone – MR208047L4	
OTI Disclaimer:	Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <u>custsupport@origene.com</u> or by calling 301.340.3188 option 3 for pricing and delivery.
	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method	<ol> <li>Centrifuge at 5,000xg for 5min.</li> <li>Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>Close the tube and incubate for 10 minutes at room temperature.</li> <li>Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM 009422.2, NP 033448.2</u>
RefSeq Size:	3044 bp
RefSeq ORF:	1506 bp
Locus ID:	22030
UniProt ID:	<u>P39429</u>
Cytogenetics:	2 A3

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### **GRIGENE** Traf2 (NM\_009422) Mouse Tagged Lenti ORF Clone – MR208047L4

# Gene Summary:Regulates activation of NF-kappa-B and JNK and plays a central role in the regulation of cell<br/>survival and apoptosis. Required for normal antibody isotype switching from IgM to IgG. Has<br/>E3 ubiquitin-protein ligase activity and promotes 'Lys-63'-linked ubiquitination of target<br/>proteins, such as BIRC3, RIPK1 and TICAM1. Is an essential constituent of several E3 ubiquitin-<br/>protein ligase complexes, where it promotes the ubiquitination of target proteins by bringing<br/>them into contact with other E3 ubiquitin ligases. Regulates BIRC2 and BIRC3 protein levels by<br/>inhibiting their autoubiquitination and subsequent degradation; this does not depend on the<br/>TRAF2 RING-type zinc finger domain. Isoform 2 does not seem to mediate activation of NF-<br/>kappa-B but inhibits isoform 1 activity. Plays a role in mediating activation of IKBKE.<br/>[UniProtKB/Swiss-Prot Function]

## **Product images:**



Circular map for MR208047L4

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US