

Product datasheet for **MR209534**

Grhl2 (NM_026496) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Grhl2 (NM_026496) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Grhl2
Synonyms:	0610015A08Rik; BOM; clft3; Tcfcp2l3
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

ORF Nucleotide Sequence:

>MR209534 representing NM_026496
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGTCACAAGAGTCGGACAATAATAAAAGACTAGTGGCCTTAGTGCCCATGCCAGTGACCTCCCTTCA
 ACACCCGAAGAGCTACACAAGTGAGGATGAGGCTGGAAGTCATATCTGGAGAACCCCTGACTGCGGC
 CACCAAGGCGATGATGAGCATCAACGGGACGAGGACAGTGCTGCCGCCCTGGGCCTGCTCTATGACTAC
 TACAAGTTCTCGAGACAAGAGACTTCTGTCTGTGAGCAAAGCAAGTGACAGCCAAGAAGACCAGGATA
 AAAGAACTGCCTTGGCACCAGTGAAGCCAGATCAATTTGAGCGGAGGCGAGAACAGAGTGCAGGTTCT
 GAAGACTGTCCCGGTGAACCTCTGTCTAAGTCAAGACCACATGGAGAATTCGAAGCGGAGCAGTACAGT
 GTATCCATCACCGAGAGCTCTGCCGTATCCCCGTGTGAGGCATCACCGTGGTAAAGCCGAGGATTTCA
 CACCGGTGTTTATGGCGCCCCGGTGCCTATCCCCGCGGACAGTGAAGGACGAGCGGTTATCTT
 TGAACAGACTCAGTACGACTGCCCTCCATAGCCAGCCACAGCTCCTATCTCAAGGACGACGAGCGCAGC
 ACGCCGGACAGCACCTACAGCGAGAGCTTAAAGGACGGCCCTCGGAGAAATTCGGAGTACTTCTGTTG
 GTGCTGACGAGTATACATATGACCAGACGGGAAGTGGTACATTTCAAGTACACCCTGGAAGCCACCAATC
 TCTCCGTGAGAAGCAGGGGGAGGGCCCCATGACCTACCTCAACAAAGGACAATTCATGCCATAACACTC
 AGTGAGACTGGAGACAACAAATGCTTCCGACACCCCATCAGCAAAGTCAAGGAGTGGTGGTGGTGGTCT
 TTAGTGAAGACAAAAACCGAGATGAGCAGCTGAAATACTGGAAGTACTGGCACTCCCGGCAGCACACTGC
 CAAGCAGAGGGTCTTGACATTGCTGATTACAAGGAGAGCTTCAACACCATCGGGAACATTGAAGAGATC
 GCATACAATGCTGTTTCTTACCTGGGATGTGAACGAGGAGGCAAGATTTTTATCACCGTGAATTGCC
 TGAGTACAGATTTCTCTCCAAAAGGGTGTAAAAGGACTTCCCTGATGATTGATGATGATGATGATGATG
 CTACAACAACCGCAGCAATAAACCCATCCACAGAGCATACTGCCAGATCAAGGTCTTCTGTGACAAGGGA
 GCAGAAAGAAAAATCCGGGATGAAGAGAGAAAGCAGAACAGAAAGAAAGGAAAGGCCAGGCCTCTCAAG
 CCCAGTGAACAACCTCTGATGGGAAGATGGCCGCCATACCGTTACAGAAGAAGAGTACATCACGTA
 CTTCAAAACCATGCCGACCTGCACTCACAGCCTGTGCTTTCATACCAGATGTTCACTTTGCAACCTA
 CAGAGGACCGGACAGGTTTATTACAACACAGACGATGAGCGAGAAGGCAGCAGCGTCTTGTAAAGCGGA
 TGTTCAAGCCCATGGAAGAGGAGTTTGGTCCAACACCGTCTAAGCAGATCAAAGAAGAAAACGTAACCG
 AGTGCTTTTATATGTGAGGAAGGAGAACGATGACGCTTCGATGCTCTGATGCTGAAATCACCCACGGTG
 AAGGGTCTGATGGAAGCGCTGTCTGAGAAGTATGGGCTGCCAGTGAGAAATCACAAGCTTTATAAGA
 AGAGCAAAAAGGCATCCTGGTCAACATGGATGACAACATCATTGAGCACTATTCAAATGAGGACACCTT
 CATCTCAACATGGAGAGCATGGTGAAGGCTTCAAGATCACGCTGATGGAGATC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence:

>MR209534 representing NM_026496
 Red=Cloning site Green=Tags(s)

MSQESDNNKRLVALVPMPSDPPFNTRRAYTSEDEAWKSYLENPLTAATKAMMSINGDEDSAAAALGLLYDY
 YKVPDRKRLLSVSKASDSQEDQDKRNLGTSEAQINLSGGENRVQVLKTPVNLCLSQDHMENSKREQYS
 VSITESSAVIPVSGITVVKAEDFTPVFMAPPVHYPRADSEEQRVVIFEQTQYDLPSIASHSSYLKDDQRS
 TPDSTYSESFKDGASEKFRSTSVGADEYTYDQTSQGFQYTLKSLRQKQEGPMTYLNKGQFYAITL
 SETGDNKCFRHPISKVRSVVMVVFSEDKNRDEQLKYWKYWHSRQHTAKQRVLDIADYKESFNITGNIIEI
 AYNAVVSFTWDVNEEAKIFITVNCLSTDFSSQKGVKGLPLMIQIDTYSYNNRNSNKP.IHRAYCQIKVFCDKG
 AERKIRDEERKQNRKKGKQASQAQCNSSDGKMAAIPLQKKSIDITYFKTMPDLHSQPVLFIPDVFHANL
 QRTGQVYYNTDDEREGSSVLVKRMFRPMEEEFGPTPSKQIKEENVKRVLLYVRKENDDVFDALMLKSPTV
 KGLMEALSEKYGLPVEKITKLYKSKKILVNMDNIIIEHYSNEDTFILNEMSMVEGFKITLMEI

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_026496

ORF Size: 1875 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_026496.4](#)

RefSeq Size: 4751 bp

RefSeq ORF: 1878 bp

Locus ID: 252973


UniProt ID: [Q8K5C0](#)

Cytogenetics: 15 B3.1

MW: 71.6 kDa

Gene Summary: Transcription factor playing an important role in primary neurulation and in epithelial development. Binds directly to the consensus DNA sequence 5'-AACCGGT-3' acting as an activator and repressor on distinct target genes (PubMed:22696678). During embryogenesis, plays unique and cooperative roles with GRHL3 in establishing distinct zones of primary neurulation. Essential for closure 3 (rostral end of the forebrain), functions cooperatively with GRHL3 in closure 2 (forebrain/midbrain boundary) and posterior neuropore closure (PubMed:20654612). Regulates epithelial morphogenesis acting as a target gene-associated transcriptional activator of apical junctional complex components. Up-regulates of CLDN3 and CLDN4, as well as of RAB25, which increases the CLDN4 protein and its localization at tight junctions (PubMed:22696678). Comprises an essential component of the transcriptional machinery that establishes appropriate expression levels of CLDN4 and CDH1 in different types of epithelia (PubMed:20978075). Exhibits functional redundancy with GRHL3 in epidermal morphogenetic events such as eyelid fusion and epidermal wound repair (PubMed:21081122). In lung, forms a regulatory loop with NKX2-1 that coordinates lung epithelial cell morphogenesis and differentiation (PubMed:22955271). In keratinocytes, plays a role in telomerase activation during cellular proliferation, regulates TERT expression by binding to TERT promoter region and inhibiting DNA methylation at the 5'-CpG island, possibly by interfering with DNMT1 enzyme activity. In addition, impairs keratinocyte differentiation and epidermal function by inhibiting the expression of genes clustered at the epidermal differentiation complex (EDC) as well as GRHL1 and GRHL3 through epigenetic mechanisms (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR209534