

## **Product datasheet for MR226160L1V**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Egfr (NM\_207655) Mouse Tagged ORF Clone Lentiviral Particle

## **Product data:**

**Product Type:** Lentiviral Particles

**Product Name:** Egfr (NM\_207655) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Egfr

**Synonyms:** 9030024J15Rik; AI552599; Erbb; Errb1; Errp; wa-2; wa2; Wa5

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 207655

ORF Size: 3630 bp

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(MR226160).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 207655.2, NP 997538.1

 RefSeq Size:
 5983 bp

 RefSeq ORF:
 3633 bp

 Locus ID:
 13649

 UniProt ID:
 Q01279

Cytogenetics: 11 9.41 cM







**Gene Summary:** 

Receptor tyrosine kinase binding ligands of the EGF family and activating several signaling cascades to convert extracellular cues into appropriate cellular responses (PubMed:8404850). Known ligands include EGF, TGFA/TGF-alpha, AREG, epigen/EPGN, BTC/betacellulin, epiregulin/EREG and HBEGF/heparin-binding EGF. Ligand binding triggers receptor homoand/or heterodimerization and autophosphorylation on key cytoplasmic residues. The phosphorylated receptor recruits adapter proteins like GRB2 which in turn activates complex downstream signaling cascades. Activates at least 4 major downstream signaling cascades including the RAS-RAF-MEK-ERK, PI3 kinase-AKT, PLCgamma-PKC and STATs modules. May also activate the NF-kappa-B signaling cascade. Also directly phosphorylates other proteins like RGS16, activating its GTPase activity and probably coupling the EGF receptor signaling to the G protein-coupled receptor signaling. Also phosphorylates MUC1 and increases its interaction with SRC and CTNNB1/beta-catenin (By similarity). Plays a role in enhancing learning and memory performance (PubMed:20639532).[UniProtKB/Swiss-Prot Function]