

Product datasheet for MR226479L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Fan1 (NM_177893) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Fan1 (NM_177893) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Fan1

Synonyms: 6030441H18Rik; mFAN1; Mtmr15

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM_177893

ORF Size: 3060 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(MR226479).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 177893.3

 RefSeq Size:
 3475 bp

 RefSeq ORF:
 3063 bp

 Locus ID:
 330554

 UniProt ID:
 Q69ZT1

Cytogenetics: 7 C





Gene Summary:

Nuclease required for the repair of DNA interstrand cross-links (ICL) recruited at sites of DNA damage by monoubiquitinated FANCD2. Specifically involved in repair of ICL-induced DNA breaks by being required for efficient homologous recombination, probably in the resolution of homologous recombination intermediates (By similarity). Not involved in DNA double-strand breaks resection. Acts as a 5'-3' exonuclease that anchors at a cut end of DNA and cleaves DNA successively at every third nucleotide, allowing to excise an ICL from one strand through flanking incisions (PubMed:24981866). Probably keeps excising with 3'-flap annealing until it reaches and unhooks the ICL. Acts at sites that have a 5'-terminal phosphate anchor at a nick or a 1- or 2-nucleotide flap and is augmented by a 3' flap (By similarity). Also has endonuclease activity toward 5'-flaps (PubMed:24981866). [UniProtKB/Swiss-Prot Function]