

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for MR226857L4V

Tlr1 (NM_030682) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Tlr1 (NM_030682) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Tlr1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_030682
ORF Size:	2385 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR226857).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 030682.2, NP 109607.1</u>
RefSeq Size:	2721 bp
RefSeq ORF:	2388 bp
Locus ID:	21897
UniProt ID:	Q9EPQ1
Cytogenetics:	5 33.53 cM



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Gene Summary:Participates in the innate immune response to microbial agents. Specifically recognizes
diacylated and triacylated lipopeptides. Cooperates with TLR2 to mediate the innate immune
response to bacterial lipoproteins or lipopeptides. Forms the activation cluster
TLR2:TLR1:CD14 in response to triacylated lipopeptides, this cluster triggers signaling from
the cell surface and subsequently is targeted to the Golgi in a lipid-raft dependent pathway.
Acts via MYD88 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the
inflammatory response (By similarity). Acts as a coreceptor for M.tuberculosis lipoproteins
LprG, LpqH and PhoS1 (pstS1), in conjunction with TLR2 and for some but not all lipoproteins
CD14 and/or CD36. The lipoproteins act as agonists to modulate antigen presenting cell
functions in response to the pathogen (PubMed:19362712).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US