

Product datasheet for PH312067

PML Protein (PML) (NM_033239) Human Mass Spec Standard

Product data:

Product Type:	Mass Spec Standards
Description:	PML MS Standard C13 and N15-labeled recombinant protein (NP_150242)
Species:	Human
Expression Host:	HEK293
Expression cDNA Clone or AA Sequence:	RC212067
Predicted MW:	90.5 kDa
Protein Sequence:	>RC212067 representing NM_033239 Red=Cloning site Green=Tags(s)
	MEPAPARSPRPQQDPARPQEPMPPEPTSEGRQSPSPSPSPTERAPASEEEFQFLRCQQCQAEAKCPKLL PCLHTLCSGCLASGMQCPICQAPWPLGADTPALDNVFFESLQRRLSVYRQIVDAQAVCTRCKESADFWC FECEQLLCAKCFEAHQWFLKHEARPLAELRNQSVREFLDGTRKTNNIFCSNPNHRTPTLTSIYCRGCSKP LCCSCALLDSSHSELKCDISAEIQQRQEELDAMTQALQEQDSAFGAVHAQMHAAVGQLGRARAETEELIR ERVVRQVVAHVRAQERELLEAVDARYQRDYEMASRLGRLDAVLQRIRTGSALVQRMKCYASDQEVLDMHG FLRQALCRLRQEEPOSLQAAVRTDGFDEFKVLQDLSSCITQGKDAAVSKKASPEAASTPRDPIDVDLPE EAERVKAQVQALGLAEAQPMVVQSVPGAHPVPVYAFSFKGSPYGEDVSNNTTAAQKRKCSQTQCPRKVIK MESEEGKEARLARSSPEQPRPSTKAVSPPHLDGPPSPRSPVIGSEVFLPNSNHVASGAGEAEERVVVIS SSESDAENSCMEPMETAEQSSPAHSSPAHSSPAHSSPVQSLLRAGASSLPCGTYHPPAWPPHPAEQ AATPDAEPHSEPPDHQERPAVHRGIRYLLYRAQRAIRLRHALRHPQLHRAPIRTWSPHVQASTPAITG PLNHPANAQEHPAQLQRGISPPHRIRGAVRSRSLRGSLSLQWLNFFALPFSMASQLDMSSVVGAG ESRAQTLGAGVPPGDSVRSMEASQVQVPLEASPIITFPPPCAPERPPISPVPGARQAGL
	SGPTRTRRLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Labeling Method:	Labeled with [U- ¹³ C ₆ , ¹⁵ N ₄]-L-Arginine and [U- ¹³ C ₆ , ¹⁵ N ₂]-L-Lysine
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3
Storage:	Store at -80°C. Avoid repeated freeze-thaw cycles.
Stability:	Stable for 3 months from receipt of products under proper storage and handling conditions.
RefSeq:	NP_150242



[View online »](#)


RefSeq Size:	3088
RefSeq ORF:	2487
Synonyms:	MYL; PP8675; RNF71; TRIM19
Locus ID:	5371
UniProt ID:	P29590
Cytogenetics:	15q24.1

Summary: The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. This phosphoprotein localizes to nuclear bodies where it functions as a transcription factor and tumor suppressor. Its expression is cell-cycle related and it regulates the p53 response to oncogenic signals. The gene is often involved in the translocation with the retinoic acid receptor alpha gene associated with acute promyelocytic leukemia (APL). Extensive alternative splicing of this gene results in several variations of the protein's central and C-terminal regions; all variants encode the same N-terminus. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2008]

Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: Acute myeloid leukemia, Pathways in cancer, Ubiquitin mediated proteolysis

Product images:



Coomassie blue staining of purified PML protein (Cat# [TP312067]). The protein was produced from HEK293T cells transfected with PML cDNA clone (Cat# [RC212067]) using MegaTran 2.0 (Cat# [TT210002]).