

Product datasheet for PH312425

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

AMHR2 (NM_020547) Human Mass Spec Standard

Product data:

Product Type: Mass Spec Standards

Description: AMHR2 MS Standard C13 and N15-labeled recombinant protein (NP_065434)

Species:HumanExpression Host:HEK293

Expression cDNA Clone

RC212425

or AA Sequence:

Protein Sequence:

Predicted MW: 62.6 kDa

>RC212425 representing NM_020547 Red=Cloning site Green=Tags(s)

MLGSLGLWALLPTAVEAPPNRRTCVFFEAPGVRGSTKTLGELLDTGTELPRAIRCLYSRCCFGIWNLTQD RAQVEMQGCRDSDEPGCESLHCDPSPRAHPSPGSTLFTCSCGTDFCNANYSHLPPPGSPGTPGSQGPQAA PGESIWMALVLLGLFLLLLLLGSIILALLQRKNYRVRGEPVPEPRPDSGRDWSVELQELPELCFSQQVI REGGHAVVWAGQLQGKLVAIKAFPPRSVAQFQAERALYELPGLQHDHIVRFITASRGGPGRLLSGPLLVL ELHPKGSLCHYLTQYTSDWGSSLRMALSLAQGLAFLHEERWQNGQNKPGIAHRDLSSQNVLIREDGSCAI GDLGLALVLPGLTQPPAWTPTQPQGPAAIMEAGTQRYMAPELLDKTLDLQDWGMALRRADIYSLALLLWE ILSRCPDLRPDSSPPPFQLAYEAELGNTPTSDELWALAVQERRRPYIPSTWRCFATDPDGLRELLEDCWD ADPEARLTAECVQQRLAALAHPQESHPFPESCPRGCPPLCPEDCTSIPAPTILPCRPQRSACHFSVQQGP

CSRNPQPACTLSPV

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

Purity: > 80% as determined by SDS-PAGE and Coomassie blue staining

Concentration: >0.05 μg/μL as determined by microplate BCA method

Labeling Method: Labeled with [U- 13C6, 15N4]-L-Arginine and [U- 13C6, 15N2]-L-Lysine

Buffer: 25 mM Tris-HCl, 100 mM glycine, pH 7.3

Storage: Store at -80°C. Avoid repeated freeze-thaw cycles.

Stability: Stable for 3 months from receipt of products under proper storage and handling conditions.

RefSeq: NP 065434

RefSeq Size:1855RefSeq ORF:1722



AMHR2 (NM_020547) Human Mass Spec Standard - PH312425

Synonyms: AMHR; MISR2; MISRII; MRII

Locus ID: 269

 UniProt ID:
 Q16671

 Cytogenetics:
 12q13.13

Summary: This gene encodes the receptor for the anti-Mullerian hormone (AMH) which, in addition to

testosterone, results in male sex differentiation. AMH and testosterone are produced in the testes by different cells and have different effects. Testosterone promotes the development


of male genitalia while the binding of AMH to the encoded receptor prevents the

development of the mullerian ducts into uterus and Fallopian tubes. Mutations in this gene are associated with persistent Mullerian duct syndrome type II. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Sep 2009]

Protein Families: Druggable Genome, Protein Kinase, Transmembrane

Protein Pathways: Cytokine-cytokine receptor interaction, TGF-beta signaling pathway

Product images:



Coomassie blue staining of purified AMHR2 protein (Cat# [TP312425]). The protein was produced from HEK293T cells transfected with AMHR2 cDNA clone (Cat# [RC212425]) using MegaTran 2.0 (Cat# [TT210002]).