

## Product datasheet for PH322829

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## JNK1 (MAPK8) (NM 139046) Human Mass Spec Standard

**Product data:** 

**Product Type:** Mass Spec Standards

**Description:** MAPK8 MS Standard C13 and N15-labeled recombinant protein (NP\_620634)

Species: Human **HEK293 Expression Host: Expression cDNA Clone** 

or AA Sequence:

RC222829

Predicted MW: 43.8 kDa

>RC222829 representing NM\_139046 **Protein Sequence:** 

Red=Cloning site Green=Tags(s)

MSRSKRDNNFYSVEIGDSTFTVLKRYQNLKPIGSGAQGIVCAAYDAILERNVAIKKLSRPFQNQTHAKRA YRELVLMKCVNHKNIIGLLNVFTPQKSLEEFQDVYIVMELMDANLCQVIQMELDHERMSYLLYQMLCGIK HLHSAGIIHRDLKPSNIVVKSDCTLKILDFGLARTAGTSFMMTPYVVTRYYRAPEVILGMGYKENVDIWS VGCIMGEMIKGGVLFPGTDHIDQWNKVIEQLGTPCPEFMKKLQPTVRTYVENRPKYAGYSFEKLFPDVLF PADSEHNKLKASQARDLLSKMLVIDASKRISVDEALQHPYINVWYDPSEAEAPPPKIPDKQLDEREHTIE

EWKELIYKEVMDLEERTKNGVIRGQPSPLAQVQQ

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag: C-Myc/DDK

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Concentration:** >0.05 µg/µL as determined by microplate BCA method

**Labeling Method:** Labeled with [U-13C6, 15N4]-L-Arginine and [U-13C6, 15N2]-L-Lysine

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3

Storage: Store at -80°C. Avoid repeated freeze-thaw cycles.

Stability: Stable for 3 months from receipt of products under proper storage and handling conditions.

RefSeq: NP 620634

RefSeg Size: 1417 RefSeq ORF: 1152

JNK; JNK-46; JNK1; JNK1A2; JNK21B1/2; PRKM8; SAPK1; SAPK1c Synonyms:

Locus ID: 5599





 UniProt ID:
 P45983

 Cytogenetics:
 10q11.22

**Summary:** The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as

an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and

development. This kinase is activated by various cell stimuli, and targets specific transcription factors, and thus mediates immediate-early gene expression in response to cell stimuli. The activation of this kinase by tumor-necrosis factor alpha (TNF-alpha) is found to be required for TNF-alpha induced apoptosis. This kinase is also involved in UV radiation induced apoptosis, which is thought to be related to cytochrom c-mediated cell death pathway. Studies of the mouse counterpart of this gene suggested that this kinase play a key role in T cell proliferation, apoptosis and differentiation. Several alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, Apr 2016]


**Protein Families:** Druggable Genome, ES Cell Differentiation/IPS, Protein Kinase

**Protein Pathways:** Adipocytokine signaling pathway, Colorectal cancer, Epithelial cell signaling in Helicobacter

pylori infection, ErbB signaling pathway, Fc epsilon RI signaling pathway, Focal adhesion, GnRH signaling pathway, Insulin signaling pathway, MAPK signaling pathway, Neurotrophin signaling pathway, NOD-like receptor signaling pathway, Pancreatic cancer, Pathways in cancer, Progesterone-mediated oocyte maturation, RIG-I-like receptor signaling pathway, Toll-

like receptor signaling pathway, Type II diabetes mellitus, Wnt signaling pathway

## **Product images:**



Coomassie blue staining of purified MAPK8 protein (Cat# [TP322829]). The protein was produced from HEK293T cells transfected with MAPK8 cDNA clone (Cat# [RC222829]) using MegaTran 2.0 (Cat# [TT210002]).