

## Product datasheet for **R1378T**

### Rat IgG (H+L chain, adsorbed) Goat Polyclonal Antibody

#### Product data:

<b>Product Type:</b>	Secondary Antibodies
<b>Product Name:</b>	Rat IgG (H+L chain, adsorbed) Goat Polyclonal Antibody
<b>Recommended Dilution:</b>	Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency.
<b>Reactivity:</b>	Rat
<b>Host:</b>	Goat
<b>Immunogen:</b>	Rat IgG whole molecule.
<b>Formulation:</b>	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, containing 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v) Sodium Azide as preservative. Label: TRITC State: Lyophilized purified Ig fraction. Label: Tetramethylrhodamine isothiocyanate ; Molecular Weight 444 daltons) Absorption emission: 550 nm / 570 nm Molar ratio: 2.9 moles TRITC per mole of Goat IgG.
<b>Reconstitution Method:</b>	Restore with 1.0 ml of deionized water (or equivalent).
<b>Concentration:</b>	lot specific
<b>Purification:</b>	Immunoaffinity chromatography.
<b>Conjugation:</b>	TRITC
<b>Storage:</b>	Store vial at 4°C prior to restoration. For extended storage reconstitute product with 50% glycerol instead of water and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 4°C as an undiluted liquid. Dilute only prior to immediate use. Avoid cycles of freezing and thawing.



[View online »](#)