

Product datasheet for R1390F

Rabbit IgG (H+L chain) (Fab Fragment) Goat Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Rabbit IgG (H+L chain) (Fab Fragment) Goat Polyclonal Antibody
Applications:	ELISA, FC, IF
Recommended Dilution:	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. Recommended dilutions: Flow cytometry 1:2,000 - 1:10,000 Immunofluorescence 1:1,000 - 1:5,000 FLISA:1:10,000-1:50,000.
Reactivity:	Rabbit
Host:	Goat
Immunogen:	Rabbit IgG whole molecule
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, containing 0.01% (w/v) Sodium Azide as a preservative and 10 mg/ml Bovine Serum Albumin (BSA), IgG and Protease free, as stabilizer Label: FITC State: Lyophilized Label: Fluorescein isothiocyanate (Molecular Weight 390 daltons) Absorption emission: 495 nm / 528 nm Molar ratio: 2.9 moles FITC per mole of Goat IgG Fab
Reconstitution Method:	Restore with 1.0 ml of deionized water or equivalent.
Concentration:	lot specific
Purification:	Immunoaffinity chromatography
Conjugation:	FITC



[View online »](#)

Storage:

Store vial at 4°C prior to restoration. Restore with deionized water (or equivalent); centrifuge product if not completely clear after standing at room temperature. This product is stable for four weeks at 4°C as an undiluted liquid.

For extended storage reconstitute product with 50% glycerol instead of water and then aliquot contents and freeze at -20°C or below. Avoid cycles of freezing and thawing. Dilute only prior to immediate use.