

Product datasheet for RC200480L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DHCR7 (NM_001360) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: DHCR7 (NM_001360) Human Tagged ORF Clone Lentiviral Particle

Symbol: DHCR Synonyms: SLOS

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_001360

 ORF Size:
 1425 bp

ORF Nucleotide

Sequence:

Cytogenetics:

The ORF insert of this clone is exactly the same as(RC200480).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 001360.2

 RefSeq Size:
 2665 bp

 RefSeq ORF:
 1428 bp

 Locus ID:
 1717

 UniProt ID:
 Q9UBM7

Domains: ERG4 ERG24

Protein Families: Druggable Genome, Transmembrane

11q13.4





DHCR7 (NM_001360) Human Tagged ORF Clone Lentiviral Particle - RC200480L3V

Protein Pathways: Metabolic pathways, Steroid biosynthesis

MW: 54.5 kDa

Gene Summary: This gene encodes an enzyme that removes the C(7-8) double bond in the B ring of sterols

and catalyzes the conversion of 7-dehydrocholesterol to cholesterol. This gene is ubiquitously expressed and its transmembrane protein localizes to the endoplasmic reticulum membrane and nuclear outer membrane. Mutations in this gene cause Smith-Lemli-Opitz syndrome (SLOS); a syndrome that is metabolically characterized by reduced serum cholesterol levels and elevated serum 7-dehydrocholesterol levels and phenotypically characterized by cognitive disability, facial dysmorphism, syndactyly of second and third toes, and holoprosencephaly in severe cases to minimal physical abnormalities and near-normal intelligence in mild cases. Alternative splicing results in multiple transcript variants that

encode the same protein.[provided by RefSeq, Aug 2009]