

Product datasheet for RC201168L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Beta Arrestin 2 (ARRB2) (NM_004313) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Beta Arrestin 2 (ARRB2) (NM_004313) Human Tagged ORF Clone Lentiviral Particle

Symbol: ARRB2

Synonyms: ARB2; ARR2; BARR2

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_004313 **ORF Size:** 1227 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC201168).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 004313.3

RefSeq Size: 1936 bp
RefSeq ORF: 1230 bp
Locus ID: 409

 UniProt ID:
 P32121

 Cytogenetics:
 17p13.2

Domains: arrestin

Protein Families: Druggable Genome





Beta Arrestin 2 (ARRB2) (NM_004313) Human Tagged ORF Clone Lentiviral Particle – RC201168L2V

Protein Pathways: Chemokine signaling pathway, Endocytosis, MAPK signaling pathway, Olfactory transduction

MW: 46.1 kDa

Gene Summary: Members of arrestin/beta-arrestin protein family are thought to participate in agonist-

mediated desensitization of G-protein-coupled receptors and cause specific dampening of cellular responses to stimuli such as hormones, neurotransmitters, or sensory signals. Arrestin beta 2, like arrestin beta 1, was shown to inhibit beta-adrenergic receptor function in vitro. It is expressed at high levels in the central nervous system and may play a role in the regulation of synaptic receptors. Besides the brain, a cDNA for arrestin beta 2 was isolated from thyroid gland, and thus it may also be involved in hormone-specific desensitization of TSH receptors. Multiple alternatively spliced transcript variants encoding different isoforms

have been found for this gene. [provided by RefSeq, Mar 2012]