

## Product datasheet for RC201242L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CD20 (MS4A1) (NM\_021950) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** CD20 (MS4A1) (NM\_021950) Human Tagged ORF Clone Lentiviral Particle

Symbol: CD20

Synonyms: B1; Bp35; CD20; CVID5; FMC7; LEU-16; MS4A2; S7

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 021950

ORF Size: 891 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC201242).

OTI Disclaimer:

Sequence:

r: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 021950.3, NP 068769.2


RefSeq Size: 3331 bp
RefSeq ORF: 894 bp
Locus ID: 931
UniProt ID: P11836

Cytogenetics: 11q12.2

Domains: CD20

**Protein Families:** Druggable Genome, Transmembrane





**Protein Pathways:** Hematopoietic cell lineage

MW: 33.1 kDa

ORIGENE

**Gene Summary:** This gene encodes a member of the membrane-spanning 4A gene family. Members of this

nascent protein family are characterized by common structural features and similar

intron/exon splice boundaries and display unique expression patterns among hematopoietic cells and nonlymphoid tissues. This gene encodes a B-lymphocyte surface molecule which plays a role in the development and differentiation of B-cells into plasma cells. This family member is localized to 11q12, among a cluster of family members. Alternative splicing of this gene results in two transcript variants which encode the same protein. [provided by RefSeq,

Jul 2008]