

Product datasheet for RC201992L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GRO alpha (CXCL1) (NM 001511) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: GRO alpha (CXCL1) (NM 001511) Human Tagged ORF Clone Lentiviral Particle

Symbol: CXCL1

Synonyms: FSP; GRO1; GROa; MGSA; MGSA-a; NAP-3; SCYB1

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_001511

ORF Size: 321 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC201992).

Sequence:
OTI Disclaimer:

Domains:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001511.1</u>

 RefSeq Size:
 1184 bp

 RefSeq ORF:
 324 bp

 Locus ID:
 2919

 UniProt ID:
 P09341

 Cytogenetics:
 4q13.3

Protein Families: Druggable Genome, Secreted Protein

IL8





GRO alpha (CXCL1) (NM_001511) Human Tagged ORF Clone Lentiviral Particle - RC201992L2V

Protein Pathways: Chemokine signaling pathway, Cytokine-cytokine receptor interaction, Epithelial cell signaling

in Helicobacter pylori infection, NOD-like receptor signaling pathway

MW: 11.3 kDa

Gene Summary: This antimicrobial gene encodes a member of the CXC subfamily of chemokines. The

encoded protein is a secreted growth factor that signals through the G-protein coupled receptor, CXC receptor 2. This protein plays a role in inflammation and as a chemoattractant for neutrophils. Aberrant expression of this protein is associated with the growth and progression of certain tumors. A naturally occurring processed form of this protein has increased chemotactic activity. Alternate splicing results in coding and non-coding variants of this gene. A pseudogene of this gene is found on chromosome 4. [provided by RefSeq, Sep

2014]