

## Product datasheet for RC202461L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## HIF-1 alpha (HIF1A) (NM\_001530) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: HIF-1 alpha (HIF1A) (NM 001530) Human Tagged ORF Clone Lentiviral Particle

Symbol: HIF-1 alpha

Synonyms: bHLHe78; HIF-1-alpha; HIF-1A; HIF-1alpha; HIF1; HIF1-ALPHA; MOP1; PASD8

Mammalian Cell

Selection:

ACCN:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

NM 001530

Tag: Myc-DDK

ORF Size: 2478 bp

**ORF Nucleotide** 

leotide The ORE

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(RC202461).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through paturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 001530.2

RefSeq Size: 3958 bp
RefSeq ORF: 2481 bp
Locus ID: 3091

UniProt ID: Q16665
Cytogenetics: 14q23.2

**Domains:** PAS, HLH, PAC

**Protein Families:** Transcription Factors





**Protein Pathways:** mTOR signaling pathway, Pathways in cancer, Renal cell carcinoma

**MW:** 92.5 kDa

Gene Summary: This gene encodes the alpha subunit of transcription factor hypoxia-inducible factor-1 (HIF-1),

which is a heterodimer composed of an alpha and a beta subunit. HIF-1 functions as a master regulator of cellular and systemic homeostatic response to hypoxia by activating transcription of many genes, including those involved in energy metabolism, angiogenesis, apoptosis, and other genes whose protein products increase oxygen delivery or facilitate

metabolic adaptation to hypoxia. HIF-1 thus plays an essential role in embryonic

vascularization, tumor angiogenesis and pathophysiology of ischemic disease. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene.

[provided by RefSeq, Jul 2011]