

Product datasheet for RC204872L1

XPA (NM_000380) Human Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	XPA (NM_000380) Human Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	XPA
Synonyms:	XP1; XPAC
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC204872).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.


ACCN:	NM_000380
ORF Size:	819 bp




[View online >](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_000380.2
RefSeq Size:	1491 bp
RefSeq ORF:	822 bp
Locus ID:	7507
UniProt ID:	P23025
Cytogenetics:	9q22.33
Domains:	XPA_C
Protein Families:	Druggable Genome
Protein Pathways:	Nucleotide excision repair
MW:	31.4 kDa
Gene Summary:	This gene encodes a zinc finger protein plays a central role in nucleotide excision repair (NER), a specialized type of DNA repair. NER is responsible for repair of UV radiation-induced photoproducts and DNA adducts induced by chemical carcinogens and chemotherapeutic drugs. The encoded protein interacts with DNA and several NER proteins, acting as a scaffold to assemble the NER incision complex at sites of DNA damage. Mutations in this gene cause Xeroderma pigmentosum complementation group A (XP-A), an autosomal recessive skin disorder featuring hypersensitivity to sunlight and increased risk for skin cancer. [provided by RefSeq, Aug 2017]

Product images:



Circular map for RC204872L1



Double digestion of RC204872L1 using SgfI and MluI