

Product datasheet for RC206897L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Ubiquilin (UBQLN1) (NM_013438) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Ubiquilin (UBQLN1) (NM_013438) Human Tagged ORF Clone Lentiviral Particle

Symbol: Ubiquilin

Synonyms: DA41; DSK2; PLIC-1; UBQN; XDRP1

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_013438 **ORF Size:** 1767 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC206897).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 013438.3</u>

RefSeq Size: 4177 bp
RefSeq ORF: 1770 bp
Locus ID: 29979
UniProt ID: Q9UMX0

Cytogenetics: 9q21.2-q21.3 **Domains:** UBA, UBQ, STI1

Protein Families: Druggable Genome





MW:

62.6 kDa

Gene Summary:

This gene encodes an ubiquitin-like protein (ubiquilin) that shares a high degree of similarity with related products in yeast, rat and frog. Ubiquilins contain an N-terminal ubiquitin-like domain and a C-terminal ubiquitin-associated domain. They physically associate with both proteasomes and ubiquitin ligases, and thus are thought to functionally link the ubiquitination machinery to the proteasome to affect in vivo protein degradation. This ubiquilin has also been shown to modulate accumulation of presenilin proteins, and it is found in lesions associated with Alzheimer's and Parkinson's disease. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]