

# Product datasheet for RC211042

### H2BC18 (NM\_001024599) Human Tagged ORF Clone

### **Product data:**

#### OriGene Technologies, Inc.


9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Expression Plasmids
Product Name:	H2BC18 (NM_001024599) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	H2BC18
Synonyms:	HIST2H2BF
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	<pre>&gt;RC211042 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)</pre>
	TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C
	ATGCCGGATCCAGCGAAATCCGCTCCTGCTCCCAAGAAGGGCTCCAAAAAGGCTGTTACGAAAGTGCAGA AGAAGGACGGCAAGAAGCGCAAGCGCAGCCGCAAGGAGAGAGCTACTCCGTTTACGTGTACAAGGTGCTGAA GCAGGTCCACCCCGACACCGGCATCTCGTCCAAGGCCATGGGCATCATGAACTCCTTCGTCAACGACATC TTCGAGCGCATCGCGGGAGAGGCGTCCCGCCTGGCGCACTACAACAAGCGCTCCACCATCACATCCCGCG AGATCCAGACGGCCGTGCGCGCTGCTGCCGGCGAGCTGGCCAAGCACGCCGTGTCCGAGGGCACCAA GGCGGTCACCAAGTACACCAGCTCGAAG
	ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT ACAAGGATGACGACGATAAG <b>GTTTAA</b>
Protein Sequence:	>RC211042 protein sequence Red=Cloning site Green=Tags(s)
	MPDPAKSAPAPKKGSKKAVTKVQKKDGKKRKRSRKESYSVYVYKVLKQVHPDTGISSKAMGIMNSFVNDI FERIAGEASRLAHYNKRSTITSREIQTAVRLLLPGELAKHAVSEGTKAVTKYTSSK
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Chromatograms:	https://cdn.origene.com/chromatograms/mk6378_h05.zip
<b>Restriction Sites:</b>	Sgfl-Mlul



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **Cloning Scheme:**




\* The last codon before the Stop codon of the ORF

ACCN:	NM_001024599
	_
ORF Size: OTI Disclaimer:	378 bp The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol> <li>Centrifuge at 5,000xg for 5min.</li> <li>Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>Close the tube and incubate for 10 minutes at room temperature.</li> <li>Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
RefSeq:	<u>NM 001024599.5</u>
RefSeq Size:	495 bp
RefSeq ORF:	381 bp
Locus ID:	440689
UniProt ID:	<u>Q5QNW6</u>


This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

<b>ORÏGENE</b> H2BC18 (NM_001024599) Human Tagged ORF Clone – RC211042		
Cytogenetics:	1q21.2	
Protein Pathways:	Systemic lupus erythematosus	
MW:	13.9 kDa	
Gene Summary:	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene encodes a replication-dependent histone that is a member of the histone H2B family and is found in a histone cluster on chromosome 1. [provided by RefSeq, Aug 2015]	

## **Product images:**




Circular map for RC211042



Western blot validation of overexpression lysate (Cat# [LY422516]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC211042 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Coomassie blue staining of purified HIST2H2BF protein (Cat# [TP311042]). The protein was produced from HEK293T cells transfected with HIST2H2BF cDNA clone (Cat# RC211042) using MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US