

## Product datasheet for RC213804L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Optineurin (OPTN) (NM\_001008212) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Optineurin (OPTN) (NM\_001008212) Human Tagged ORF Clone Lentiviral Particle

Symbol: Optineurin

Synonyms: ALS12; FIP2; GLC1E; HIP7; HYPL; NRP; TFIIIA-INTP

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_001008212

ORF Size: 1731 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC213804).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 001008212.1

 RefSeq Size:
 3528 bp

 RefSeq ORF:
 1734 bp

 Locus ID:
 10133

 UniProt ID:
 Q96CV9

Cytogenetics: 10p13

**Protein Families:** Druggable Genome

**MW:** 65.9 kDa





## **Gene Summary:**

This gene encodes the coiled-coil containing protein optineurin. Optineurin may play a role in normal-tension glaucoma and adult-onset primary open angle glaucoma. Optineurin interacts with adenovirus E3-14.7K protein and may utilize tumor necrosis factor-alpha or Fas-ligand pathways to mediate apoptosis, inflammation or vasoconstriction. Optineurin may also function in cellular morphogenesis and membrane trafficking, vesicle trafficking, and transcription activation through its interactions with the RAB8, huntingtin, and transcription factor IIIA proteins. Alternative splicing results in multiple transcript variants encoding the same protein. [provided by RefSeq, Jul 2008]