

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for RC214815

## H3FC (HIST1H3C) (NM\_003531) Human Tagged ORF Clone


## **Product data:**

Product Type:	Expression Plasmids
Product Name:	H3FC (HIST1H3C) (NM_003531) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	H3FC
Synonyms:	H3.1; H3/c; H3C1; H3C2; H3C4; H3C6; H3C7; H3C8; H3C10; H3C11; H3C12; H3FC; HIST1H3C
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	<pre>&gt;RC214815 representing NM_003531 Red=Cloning site Blue=ORF Green=Tags(s)</pre>
	TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C
	ATGGCTCGTACGAAGCAAACAGCTCGCAAGTCTACCGGCGGCAAAGCTCCGCGCAAGCAGCTTGCTACTA AAGCAGCCCGTAAGAGCGCTCCGGCCACCGGTGGCGTGAAGAAACCTCATCGCTACCGCCCGGGCACCGT GGCCTTGCGCGAAATCCGTCGCTACCAGAAGTCCACCGAGCTGCTGATCCGGAAGCTGCCGTTCCAGCGC CTGGTGCGAGAAATCGCCCAGGACTTCAAAACCGACCTGCGTTTCCAGAGCTCTGCGGTGATGGCGCTGC AGGAGGCTTGTGAGGCCTACCTGGTGGGACTCTTCGAAGACACCAATCTGTGCGCTATTCACGCTAAACG CGTCACCATCATGCCCAAAGATATCCAGCTGGCACGTCGCATCCGTGGGGAAAGGGCA
	AGCGGACCGACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCC TGGATTACAAGGATGACGACGATAAG <b>GTTTAA</b>
Protein Sequence:	<pre>&gt;RC214815 representing NM_003531 Red=Cloning site Green=Tags(s)</pre>
	MARTKQTARKSTGGKAPRKQLATKAARKSAPATGGVKKPHRYRPGTVALREIRRYQKSTELLIRKLPFQR LVREIAQDFKTDLRFQSSAVMALQEACEAYLVGLFEDTNLCAIHAKRVTIMPKDIQLARRIRGERA
	SGPTRTRRLEQKLISEEDLAANDILDYKDDDDKV
Chromatograms:	https://cdn.origene.com/chromatograms/mg3328_e01.zip
Restriction Sites:	Sgfl-RsrII



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **Cloning Scheme:**




ORF Size:408 bpOTI Disclaimer:The molecular sequence of this clone aligns with the genereference only. However, individual transcript sequence naturally occurring variations (e.g. polymorphisms), each clone is substantially in agreement with the reference, be variants is recommended prior to use. More infoOTI Annotation:This clone was engineered to express the complete ORF varies depending on the nature of the gene.Components:The ORF clone is ion-exchange column purified and ship containing 10ug of transfection-ready, dried plasmid DNReconstitution Method:1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile wate 3. Close the tube and incubate for 10 minutes at room t 4. Briefly vortex the tube and then do a quick spin (less the tube bottom. 5. Store the suspended plasmid at -20°C. The DNA is statishipping when stored at -20°C.RefSeq:NM 003531.3	s of the same gene can differ through n with its own valid existence. This
<ul> <li>reference only. However, individual transcript sequence naturally occurring variations (e.g. polymorphisms), ead clone is substantially in agreement with the reference, be variants is recommended prior to use. More info</li> <li>OTI Annotation: This clone was engineered to express the complete ORF varies depending on the nature of the gene.</li> <li>Components: The ORF clone is ion-exchange column purified and ship containing 10ug of transfection-ready, dried plasmid DN</li> <li>Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile wate 3. Close the tube and incubate for 10 minutes at room t 4. Briefly vortex the tube and then do a quick spin (less the tube bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is statishipping when stored at -20°C.</li> </ul>	s of the same gene can differ through n with its own valid existence. This
<ul> <li>varies depending on the nature of the gene.</li> <li>Components: The ORF clone is ion-exchange column purified and ship containing 10ug of transfection-ready, dried plasmid DN</li> <li>Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile wate</li> <li>3. Close the tube and incubate for 10 minutes at room t</li> <li>4. Briefly vortex the tube and then do a quick spin (less to at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is statistical shipping when stored at -20°C.</li> </ul>	
<ul> <li>containing 10ug of transfection-ready, dried plasmid DN</li> <li>Reconstitution Method:         <ol> <li>Centrifuge at 5,000xg for 5min.</li> <li>Carefully open the tube and add 100ul of sterile wate</li> <li>Close the tube and incubate for 10 minutes at room t</li> <li>Briefly vortex the tube and then do a quick spin (less tat the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is stat shipping when stored at -20°C.</li> </ol> </li> </ul>	with an expression tag. Expression
<ol> <li>Carefully open the tube and add 100ul of sterile wate</li> <li>Close the tube and incubate for 10 minutes at room t</li> <li>Briefly vortex the tube and then do a quick spin (less at the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is statistical shipping when stored at -20°C.</li> </ol>	•
<b>RefSeq:</b> <u>NM 003531.3</u>	emperature. han 5000xg) to concentrate the liquid
RefSeq Size: 459 bp	
RefSeq ORF: 411 bp	
Locus ID: 8352	
UniProt ID: <u>P68431</u>	


This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Service Marc (HIST1H3C) (NM_003531) Human Tagged ORF Clone – RC214815		
Cytogenetics:	6p22.2	
Protein Pathways:	Systemic lupus erythematosus	
MW:	15.2 kDa	
Gene Summary:	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6. [provided by RefSeq, Aug 2015]	

## **Product images:**



Circular map for RC214815



Western blot validation of overexpression lysate (Cat# [LY418619]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC214815 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US