

Product datasheet for RC216866

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

H3C14 (NM 021059) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: H3C14 (NM_021059) Human Tagged ORF Clone

Tag: Myc-DDK Symbol: H3C14

Synonyms: H3; H3.2; H3/M; H3C13; H3C15; H3F2; H3FM; H3FN; HIST2H3C

Mammalian Cell No

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)ORF Nucleotide>RC216866 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC216866 protein sequence

Red=Cloning site Green=Tags(s)

MARTKQTARKSTGGKAPRKQLATKAARKSAPATGGVKKPHRYRPGTVALREIRRYQKSTELLIRKLPFQR LVREIAQDFKTDLRFQSSAVMALQEASEAYLVGLFEDTNLCAIHAKRVTIMPKDIQLARRIRGERA

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6473 d07.zip


Restriction Sites: Sgfl-Mlul





Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_021059

ORF Size: 408 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: NM 021059.3

 RefSeq Size:
 507 bp

 RefSeq ORF:
 411 bp

 Locus ID:
 126961

 UniProt ID:
 Q71DI3



Cytogenetics: 1q21.2


Protein Pathways: Systemic lupus erythematosus

MW: 15.4 kDa


Gene Summary:

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in a histone cluster on chromosome 1. This gene is one of four histone genes in the cluster that are duplicated; this record represents the telomeric copy. [provided by RefSeq, Aug 2015]

Product images:



Circular map for RC216866



Western blot validation of overexpression lysate (Cat# [LY412125]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC216866 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).