

Product datasheet for RC217003L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MET (NM_000245) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: MET (NM_000245) Human Tagged ORF Clone Lentiviral Particle

Symbol: MET

Synonyms: AUTS9; c-Met; DFNB97; HGFR; RCCP2

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_000245 **ORF Size:** 4170 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC217003).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 000245.2

 RefSeq Size:
 6641 bp

 RefSeq ORF:
 4173 bp

 Locus ID:
 4233

 UniProt ID:
 P08581

 Cytogenetics:
 7q31.2

Protein Families: Druggable Genome, Protein Kinase, Transmembrane





MET (NM_000245) Human Tagged ORF Clone Lentiviral Particle - RC217003L2V

Protein Pathways: Adherens junction, Axon guidance, Colorectal cancer, Cytokine-cytokine receptor interaction,

Endocytosis, Epithelial cell signaling in Helicobacter pylori infection, Focal adhesion,

Melanoma, Pathways in cancer, Renal cell carcinoma

MW: 155.54 kDa

Gene Summary: This gene encodes a member of the receptor tyrosine kinase family of proteins and the

product of the proto-oncogene MET. The encoded preproprotein is proteolytically processed to generate alpha and beta subunits that are linked via disulfide bonds to form the mature receptor. Further processing of the beta subunit results in the formation of the M10 peptide, which has been shown to reduce lung fibrosis. Binding of its ligand, hepatocyte growth factor, induces dimerization and activation of the receptor, which plays a role in cellular survival, embryogenesis, and cellular migration and invasion. Mutations in this gene are associated with papillary renal cell carcinoma, hepatocellular carcinoma, and various head and neck cancers. Amplification and overexpression of this gene are also associated with multiple

human cancers. [provided by RefSeq, May 2016]