

OriGene Technologies, Inc.


9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC221271L4

Wilms Tumor Protein (WT1) (NM_024424) Human Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Wilms Tumor Protein (WT1) (NM_024424) Human Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	Wilms Tumor Protein
Synonyms:	AWT1; GUD; NPHS4; WAGR; WIT-2; WT33
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC221271).
Restriction Sites:	Sgfl-Mlul
Cloning Scheme:	
	Cloning sites used for ORF Shuttling:
	Sgf I ORF Miu I GCG ATC GCC ATG // NNN ACG CGT



ACCN: ORF Size: NM_024424 1542 bp



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Vilms Tumor Protein (WT1) (NM_024424) Human Tagged Lenti ORF Clone – RC221271L4
OTI Disclaimer:	Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <u>custsupport@origene.com</u> or by calling 301.340.3188 option 3 for pricing and delivery.
	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Me	 thod: 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM 024424.2</u>
RefSeq Size:	3020 bp
RefSeq ORF:	1560 bp
Locus ID:	7490
UniProt ID:	<u>P19544</u>
Cytogenetics:	11p13
Protein Families:	Druggable Genome, Transcription Factors
MW:	55.8 kDa


~ \$%~

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Scrigene Wilms Tumor Protein (WT1) (NM_024424) Human Tagged Lenti ORF Clone – RC221271L4 Wilms Tumor Protein (WT1) (NM_024424) Human Tagged Lenti ORF Clone – RC221271L4

Gene Summary:This gene encodes a transcription factor that contains four zinc-finger motifs at the C-
terminus and a proline/glutamine-rich DNA-binding domain at the N-terminus. It has an
essential role in the normal development of the urogenital system, and it is mutated in a
small subset of patients with Wilms tumor. This gene exhibits complex tissue-specific and
polymorphic imprinting pattern, with biallelic, and monoallelic expression from the maternal
and paternal alleles in different tissues. Multiple transcript variants have been described. In
several variants, there is evidence for the use of a non-AUG (CUG) translation initiation codon
upstream of, and in-frame with the first AUG. Authors of PMID:7926762 also provide evidence
that WT1 mRNA undergoes RNA editing in human and rat, and that this process is tissue-
restricted and developmentally regulated. [provided by RefSeq, Mar 2015]

Product images:



Circular map for RC221271L4

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US