

## **Product datasheet for RC221674**

## BID (NM\_197967) Human Tagged ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** BID (NM 197967) Human Tagged ORF Clone

Tag: Myc-DDK

Symbol: BID

Synonyms: FP497

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

ORF Nucleotide >RC221674 representing NM\_197967

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGGACCGTAGCATCCCTCCGGGCCTGGTGAACGGCCTGGCCCTGCAGCTCAGGAACACCAGCCGGTCGGAGGAGGACCGGAACAGGGACCTGGCCACTGCCCTGGAGCAGCTGCTGCAGGCCTACCCTAGAGACATGGAGAAGAAGAACATGCTGGTGCTGGCCCTGCTGCTGGCCAAGAAGGTGGCCAGTCACACGCCGTCCTTGCTCGGTGATGTCTTTCACACAACAGAGGAGAATTTTATTAACCAGAACCTACGCACCTACGTGAGGAGCTTAG

CCAGAAATGGGATGGAC

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:** >RC221674 representing NM\_197967

Red=Cloning site Green=Tags(s)

MDRSIPPGLVNGLALQLRNTSRSEEDRNRDLATALEQLLQAYPRDMEKEKTMLVLALLLAKKVASHTPSL

LRDVFHTTVNFINQNLRTYVRSLARNGMD

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

**Restriction Sites:** Sgfl-Mlul




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Cloning Scheme:**





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_197967

ORF Size: 297 bp

**OTI Disclaimer:** Due to the inherent nature of this plasmid, standard methods to replicate additional amounts

of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:customercom">customercom</a> or by

calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** <u>NM 197967.2</u>

 RefSeq Size:
 2144 bp

 RefSeq ORF:
 300 bp

 Locus ID:
 637

 UniProt ID:
 P55957

Cytogenetics: <u>P55957</u>
22q11.21

**Protein Families:** Druggable Genome

Protein Pathways: Alzheimer's disease, Amyotrophic lateral sclerosis (ALS), Apoptosis, Natural killer cell

mediated cytotoxicity, p53 signaling pathway, Pathways in cancer, Viral myocarditis

**MW:** 11.1 kDa

**Gene Summary:** This gene encodes a death agonist that heterodimerizes with either agonist BAX or antagonist


BCL2, and thus regulate apoptosis. The encoded protein is a member of the BCL-2 family of cell death regulators. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8);

CASP8 cleaves this encoded protein, and the COOH-terminal part translocates to

mitochondria where it triggers cytochrome c release. Multiple alternatively spliced transcript


variants have been found. [provided by RefSeq, Aug 2020]

## **Product images:**



Circular map for RC221674





Western blot validation of overexpression lysate (Cat# [LY405092]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC221674 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).