

Product datasheet for RC221792L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Optineurin (OPTN) (NM 021980) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Optineurin (OPTN) (NM_021980) Human Tagged ORF Clone Lentiviral Particle

Symbol: Optineurin

Synonyms: ALS12; FIP2; GLC1E; HIP7; HYPL; NRP; TFIIIA-INTP

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_021980 **ORF Size:** 1731 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC221792).

OTI Disclaimer:

Sequence:

Domains:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 021980.4, NP 068815.2

zf-C2H2

RefSeq Size: 3376 bp
RefSeq ORF: 1734 bp
Locus ID: 10133
UniProt ID: Q96CV9
Cytogenetics: 10p13

Protein Families: Druggable Genome





ORIGENE

MW: 65.9 kDa

Gene Summary:

This gene encodes the coiled-coil containing protein optineurin. Optineurin may play a role in normal-tension glaucoma and adult-onset primary open angle glaucoma. Optineurin interacts with adenovirus E3-14.7K protein and may utilize tumor necrosis factor-alpha or Fas-ligand pathways to mediate apoptosis, inflammation or vasoconstriction. Optineurin may also function in cellular morphogenesis and membrane trafficking, vesicle trafficking, and transcription activation through its interactions with the RAB8, huntingtin, and transcription factor IIIA proteins. Alternative splicing results in multiple transcript variants encoding the same protein. [provided by RefSeq, Jul 2008]