

OriGene Technologies, Inc.


9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC226550L3

Retinoic Acid Receptor alpha (RARA) (NM_001145302) Human Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids	
Product Name:	Retinoic Acid Receptor alpha (RARA) (NM_001145302) Human Tagged Lenti ORF Clone	
Tag:	Myc-DDK	
Symbol:	Retinoic Acid Receptor alpha	
Synonyms:	NR1B1; RAR	
Mammalian Cell Selection:	Puromycin	
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)	
E. coli Selection:	Chloramphenicol (34 ug/mL)	
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC226550).	
Restriction Sites:	Sgfl-Mlul	
Cloning Scheme:		
	Cloning sites used for ORF Shuttling:	
	Sgf I ORF Mlu I GCG ATC GC ATG// NNÑ ACG CGT	



ACCN:	
ORF Size:	

View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

NM_001145302


1095 bp

PORIGENE Retinoic Acid Receptor alpha (RARA) (NM_001145302) Human Tagged Lenti ORF Clone – RC226550L3 RC226550L3	
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Met	 hod: 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM 001145302.1</u>
RefSeq ORF:	1098 bp
Locus ID:	5914
UniProt ID:	<u>P10276</u>
Cytogenetics:	17q21.2
Protein Families:	Druggable Genome, Nuclear Hormone Receptor, Transcription Factors
Protein Pathways:	Acute myeloid leukemia, Pathways in cancer
MW:	39.5 kDa
Gene Summary:	This gene represents a nuclear retinoic acid receptor. The encoded protein, retinoic acid receptor alpha, regulates transcription in a ligand-dependent manner. This gene has been implicated in regulation of development, differentiation, apoptosis, granulopoeisis, and transcription of clock genes. Translocations between this locus and several other loci have been associated with acute promyelocytic leukemia. Alternatively spliced transcript variants have been found for this locus.[provided by RefSeq, Sep 2010]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Product images:



Circular map for RC226550L3

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US