

## Product datasheet for RC230155L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## SHMT2 (NM\_001166358) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** SHMT2 (NM\_001166358) Human Tagged ORF Clone Lentiviral Particle

Symbol: SHMT2

**Synonyms:** GLYA; HEL-S-51e; NEDCASB; SHMT

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_001166358

ORF Size: 1515 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC230155).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 001166358.1</u>, <u>NP 001159830.1</u>

 RefSeq Size:
 2256 bp

 RefSeq ORF:
 1452 bp

 Locus ID:
 6472

 UniProt ID:
 P34897

 Cytogenetics:
 12q13.3

**Protein Pathways:** Cyanoamino acid metabolism, Glycine, serine and threonine metabolism, Metabolic

pathways, Methane metabolism, One carbon pool by folate







MW:

56 kDa

**Gene Summary:** 

This gene encodes the mitochondrial form of a pyridoxal phosphate-dependent enzyme that catalyzes the reversible reaction of serine and tetrahydrofolate to glycine and 5,10-methylene tetrahydrofolate. The encoded product is primarily responsible for glycine synthesis. The activity of the encoded protein has been suggested to be the primary source of intracellular glycine. The gene which encodes the cytosolic form of this enzyme is located on chromosome 17. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2009]