

Product datasheet for RC238089

Isocitrate dehydrogenase (IDH1) (NM_001282387) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Isocitrate dehydrogenase (IDH1) (NM_001282387) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Isocitrate dehydrogenase
Synonyms:	HEL-216; HEL-S-26; IDCD; IDH; IDP; IDPC; PICD
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RC238089 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGTCCAAAAAATCAGTGGCGTTCTGTGGTAGAGATGCAAGGAGATGAAATGACACGAATCATTGGG
AATTGATTAAGAGAACTCATTTTTCCCTACGTGGAATTGGATCTACATAGCTATGATTTAGGCATAGA
GAATCGTGATGCCACCAACGACCAAGTCACCAAGGATGCTGCAGAAGCTATAAGAAGCATAATGTTGGC
GTCAAATGTGCCACTATCACTCCTGATGAGAAGAGGGTTGAGGAGTTCAAGTTGAAACAAATGTGGAAT
CACCAAATGGCACCATACGAAATATTCTGGGTGGCACGGTCTTCAGAGAAGCCATTATCTGCAAAAATAT
CCCCCGCTTGTGAGTGGATGGTAAAACCTATCATCATAGGTCGTCATGCTTATGGGGATCAATACAGA
GCAACTGATTTTGTGTTCTGGCCTGGAAAAGTAGAGATAACCTACACACCAAGTGACGGAACCCAAA
AGGTGACATACCTGGTACATAACTTTGAAGAAGGTGGTGGTGTGCCATGGGGATGTATAATCAAGATAA
GTCAATTGAAGATTTTGCACACAGTTCCTTCAAATGGCTCTGTCTAAGGGTTGGCCTTTGTATCTGAGC
ACCAAAAACACTATTCTGAAGAAATATGATGGCGTTTTAAAGACATCTTTCAGGAGATATATGACAAGC
AGTACAAGTCCCAGTTTGAAGCTCAAAGATCTGGTATGAGCATAGGCTCATCGACGACATGGTGGCCCA
AGCTATGAAATCAGAGGGAGGCTTCATCTGGCCTGTAAAAACTATGATGGTGACGTGCAGTCCGACTCT
GTGGCCCAAGGGTATGGCTCTCTCGCATGATGACCAGCGTCTGGTTTGTCCAGATGGCAAGCAGTAG
AAGCAGAGTCTGCCACGGGACTGTAACCCGCTCACTACCGCATGTACCAGAAAGGACAGGAGACGTCCAC
CAATCCCATTGCTTCCATTTTGGCTGGACCAGAGGGTTAGCCCACAGAGCAAAGCTTGATAACAATAAA
GAGCTTGCTTCTTTGCAATGCTTTGGAAGAAGTCTCTATTGAGACAATTGAGGCTGGCTTCATGACCA
AGGACTTGGCTGCTTGCATTAAGGTTTACCCAATGTGCAACGTTCTGACTACTTGAATACATTTGAGTT
CATGGATAAACTTGAGAAAACCTGAAGATCAAACCTAGCTCAGGCCAACTT

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online >](#)

Protein Sequence: >RC238089 protein sequence
Red=Cloning site Green=Tags(s)

MSKKISGGSVEMQGDGMTRIIWELIKEKLIFPYVELDLHSYDLGIENRDATNDQVTKDAAEAIKKHNVG
 VKCATITPDEKRVVEEFKLMQWVSPNGTIRNLLGGTVFREAIIICKNIPRLVSGWVKPIIIGRHAYGDQYR
 ATDFVVPVPGPKVEITYTPSDGTQKVITYLVHNFEEGGVAMGMYNQDKSIEDFAHSSFQMALSKGWPLYLS
 TKNTILKKYDGRFKDIFQEIYDKQYKSQFEAQKIWYEHRLIDDMVAQAMKSEGGFIWACKNYDGDVQSDS
 VAQYGSGLGMMTSVLVCPDGKTVEAESAHGTVTRHYRMYQKQGETSTNPIASIFAWTRGLAHRAKLDNNK
 ELAFFANALEEVSIIETIEAGFMTKDLAACIKGLPNVQRSDYLNTEFMDKLGLENLKIKLAQAKL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mg4264_d09.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:



* The last codon before the Stop codon of the ORF

ACCN: NM_001282387

ORF Size: 1242 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001282387.1](#), [NP_001269316.1](#)

RefSeq Size: 2470 bp

RefSeq ORF: 1245 bp

Locus ID: 3417

UniProt ID: [O75874](#)


Cytogenetics: 2q34

Protein Pathways: Citrate cycle (TCA cycle), Glutathione metabolism, Metabolic pathways

MW: 46.7 kDa

Gene Summary: Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. Each NADP(+)-dependent isozyme is a homodimer. The protein encoded by this gene is the NADP(+)-dependent isocitrate dehydrogenase found in the cytoplasm and peroxisomes. It contains the PTS-1 peroxisomal targeting signal sequence. The presence of this enzyme in peroxisomes suggests roles in the regeneration of NADPH for intraperoxisomal reductions, such as the conversion of 2, 4-dienoyl-CoAs to 3-enoyl-CoAs, as well as in peroxisomal reactions that consume 2-oxoglutarate, namely the alpha-hydroxylation of phytanic acid. The cytoplasmic enzyme serves a significant role in cytoplasmic NADPH production. Alternatively spliced transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Sep 2013]

Product images:



Circular map for RC238089