

## Product datasheet for **RC400096**

### Isocitrate dehydrogenase (IDH1) (NM\_005896) Human Mutant ORF Clone

#### Product data:

Product Type:	Mutant ORF Clones
Product Name:	Isocitrate dehydrogenase (IDH1) (NM_005896) Human Mutant ORF Clone
Mutation Description:	R132H
Affected Codon#:	132
Affected NT#:	c. 395
Nucleotide Mutation:	IDH1 mutant (R132H), Myc-DDK-tagged ORF clone of Homo sapiens isocitrate dehydrogenase 1 (NADP+), soluble (IDH1) as transfection-ready DNA
Effect:	Missense
Symbol:	Isocitrate dehydrogenase
Synonyms:	HEL-216; HEL-S-26; IDCD; IDH; IDP; IDPC; PICD
E. coli Selection:	Kanamycin (25 ug/mL)
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
Tag:	Myc-DDK
ACCN:	NM_005896
ORF Size:	1245 bp
Restriction Sites:	Sgfl-Mlul



[View online »](#)

**ORF Nucleotide Sequence:**

>RC400096 representing NM\_005896  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGGATCGCC**

ATGTCCAAAAAATCAGTGGCGTTCTGTGGTAGAGATGCAAGGAGATGAAATGACACGAATCATTGGG  
 AATTGATTAAAGAGAACTCATTTCCTACGTGGAATTGGATCTACATAGCTATGATTTAGGCATAGA  
 GAATCGTGATGCCACCAACGACCAAGTACCAAGGATGCTGCAGAAGCTATAAAGAAGCATAATGTTGGC  
 GTCAAATGTGCCACTATCACTCCTGATGAGAAGAGGGTTGAGGAGTTCAAGTTGAAACAAATGTGGAAT  
 CACCAAATGGCACCATACGAAATATTCTGGGTGGCAGGTCTTCAGAGAAGCCATTATCTGCAAAAATAT  
 CCCCCGGCTTGAGTGGATGGTAAAACCTATCATCATAGGTCATCATGCTTATGGGGATCAATACAGA  
 GCAACTGATTTTGTTCCTGGCCTGGAAAAGTAGAGATAACCTACACACCAAGTGACGGAACCCAAA  
 AGGTGACATACCTGGTACATAACTTTGAAGAAGTGGTGGTGTGCCATGGGGATGTATAATCAAGATAA  
 GTCAATTGAAGATTTTGACACAGTTCCTTCAAATGGCTCTGTCTAAGGGTTGGCTTTGTATCTGAGC  
 ACCAAAAACACTATTCTGAAGAAATATGATGGCGTTTTAAAGACATCTTTCAGGAGATATATGACAAGC  
 AGTACAAGTCCCAGTTTGAAGCTCAAAGATCTGGTATGAGCATAGGCTCATCGACGACATGGTGGCCCA  
 AGCTATGAAATCAGAGGGAGGCTTCATCTGGCCTGTA AAAACTATGATGGTGACGTGCAGTCGGACTCT  
 GTGGCCCAAGGGTATGGCTCTCTCGGCATGATGACCAGCGTGTGGTTTGTCCAGATGGCAAGACAGTAG  
 AAGCAGAGGCTGCCACGGGACTGTAACCCGCTACTACCGCATGTACCAGAAAGGACAGGAGACGTCAC  
 CAATCCCATTGCTCCATTTTGCCTGGACCAGAGGGTTAGCCCCAGAGCAAAGCTTGATAACAATAAA  
 GAGCTTGCCTTTTGAATGCTTTGGAAGAAGTCTCTATTGAGACAATTGAGGCTGGCTTCATGACCA  
 AGGACTTGGCTGCTTGCATTAAGGTTTACCCAATGTGCAACGTTCTGACTACTTGAATACATTTGAGTT  
 CATGGATAAATTTGGAGAAAACCTGAAGATCAAACCTAGCTCAGGCCAAACTT

**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
 ACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:**

>RC400096 representing NM\_005896  
 Red=Cloning site Green=Tags(s)

MSKKISGGSVVEMQGDDEMTRIIWELIKEKLIFFPYVELDLHSYDLGIENRDATNDQVTKDAAEAIKKNHVG  
 VKCATITPDEKRVEEFKQKQMWKSPNGTIRNILGGTVFREAIICKNIPRLVSGWVKPIIIGHHAYGDQYR  
 ATDFVVPGPVKVEITYTPSDGTQKVTYLVHNFEEGGVAMGMYNQDKSIEDFAHSSFQMALSKGWPLYLS  
 TKNTILKKYDGRFKDIFQEIYDKQYKSQFEAQKIWYEHRLIDDMVAQAMKSEGGFIWACKNYDGDVQSDS  
 VAQGYGSLGMMTSVLVCPDGKTVAEAAHGTVTRHYRMYQKQETSTNPIASIFAWTRGLAHRKLDNNK  
 ELAFFANALEEVSIEETIEAGFMTKDLAACIKGLPNVQRSDYLNTFEFMDKLGENLIKLAQAKL

**TR**TRPLEQKLISEEDLAANDILDYKDDDDKV

**Chromatograms:**

/chromatograms/ja1630\_e09.zip

**Restriction Sites:**

Sgfl-Mlul

**Cloning Scheme:**

**OTI Disclaimer:**

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:**

This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:**

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**RefSeq:**

[NP\\_005887](#)

**RefSeq Size:**

2339 bp

**RefSeq ORF:**

1245 bp

**Locus ID:**

3417

**Cytogenetics:**

2q34

**Domains:**

isodh

**Protein Pathways:**

Citrate cycle (TCA cycle), Glutathione metabolism, Metabolic pathways

**MW:**

46.5 kDa

**Gene Summary:**

Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. Each NADP(+)-dependent isozyme is a homodimer. The protein encoded by this gene is the NADP(+)-dependent isocitrate dehydrogenase found in the cytoplasm and peroxisomes. It contains the PTS-1 peroxisomal targeting signal sequence. The presence of this enzyme in peroxisomes suggests roles in the regeneration of NADPH for intraperoxisomal reductions, such as the conversion of 2, 4-dienoyl-CoAs to 3-enoyl-CoAs, as well as in peroxisomal reactions that consume 2-oxoglutarate, namely the alpha-hydroxylation of phytanic acid. The cytoplasmic enzyme serves a significant role in cytoplasmic NADPH production. Alternatively spliced transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Sep 2013]